

Statistical Mechanics
E0415

Fall 2023, lecture 3
Correlations & Dissipation

“paper” groups

1. Negative temperatures Aron Dahlberg, Miklós Nemesszeghy, Emil Stråka (7.10)
2. Physics & single cell biology Eero Saariniemi, Jere Haavisto, Urho Koljonen (14.10)
3. Non-equilibrium transition & Game of Life Alisa Haukisalmi, Juuso Attenberg, Jan Loder (28.)
4. Jarzynski inequality Angelos Stathakis, Kiran Thamke, Clara Prêcheur Llarena, Tharindu Koralage (4.11)
5. Negative Representation and Instability in Democratic Elections Anna Huttunen, Emma Lehto, Heidi Kivijärvi, Atso Ikäheimo (11.)
6. Quantum phase transition Pelin Yildirim, Jonas Tjepkema, Evren Korkmazgil (18.)
7. Entropy production Ville-Eemeli Kovanen, ALgot Silvennoinen, Mikael Tuokkola (25.11)
8. Avalanches and their shape Sofia Böling, Gentry Zenuni, Valtteri Turkki (2.12)
9. Sinan Inel, (Zeno effect TBD), TBD

Take home 2


In the event that the coin is not fair, the distributions change.

If the subsequent outcomes are positively correlated, such that the possibility of going left (losing money) is increased if we previously moved left or vice versa, then the game would end faster and the distribution of steps would peak more to the smaller values for both $X_T = 0$ and $X_T = 20$.

If the subsequent outcomes are negatively correlated, such that the possibility of going left (losing money) is decreased if we previously moved left or vice versa, then the game would last longer as the amount of coins would tend to stay put leading to a distribution with higher number of steps.

2.t

Let us turn this into an exercise in gambling. You play heads and tails (toss a coin, and guess the outcome: win or lose the coin). Three questions: you start with 10 coins. Give an argument how the distribution of times it takes for you to lose all your coins looks like. What happens if you play till you have zero, or until you won all the 10 coins of your friend? Let us now consider the case where the coin is not fair: the fractional Brownian motion, where the subsequent outcomes are correlated (positively or negatively). How does that influence qualitatively those outcomes?



Comments:

The first question is actually a so-called First Passage problem. For an unbounded domain (your friend is immensely rich so you can win ad infinitum) the average time is... infinite. That is b/c the first passage time (to reach zero) τ scales with an exponent of $-3/2$ (is a power-law). You may note that this is related to the Gaussian distribution of $-1/2$ exponent, and the FP time is its derivative. "Diffusive flux".

fBm: trends and anti-trends

Correlation functions

“Fields” $s(x,y)$: how to find regularities?

$$C_t^{\text{COAR}}(\mathbf{r}) = \langle S(\mathbf{x}, t) S(\mathbf{x} + \mathbf{r}, t) \rangle$$

$$C(\mathbf{r}, \tau) = \langle S(\mathbf{x}, t) S(\mathbf{x} + \mathbf{r}, t + \tau) \rangle.$$

s : height, magnetization, activity..

Limiting behaviors ($\tau, \mathbf{x} \rightarrow \infty$)!

Scale-free behavior (2nd order phase transitions).

Check Google Maps for $s(x,y)$...
Retkeilypaikka...



Fig. 10.1 Phase separation in an Ising model, quenched (abruptly cooled) from high temperatures to zero temperature [124]. The model quickly separates into local blobs of up- and down-spins, which grow and merge, coarsening to larger blob sizes (Section 11.4.1).




Fig. 10.2 Surface annealing. An STM image of a surface, created by bombarding a close-packed gold surface with noble-gas atoms, and then allowing the irregular surface to thermally relax (Tatjana Curcic and Barbara H. Cooper [32]). The figure shows individual atomic-height steps; the arrows each show a single step pit inside another pit. The characteristic sizes of the pits and islands grow as the surface evolves and flattens.






Fig. 10.5 Power-law correlations. The schematic correlation function of figure 10.4 on a log-log plot, both at T_c (straight line, representing the power law $C \sim r^{-(d-2+\eta)}$) and above T_c (where the dependence shifts to $C \sim e^{-r/\xi(T)}$ at distances beyond the correlation length $\xi(T)$). See Chapter 12.



Experimental measures

X-rays, neutrons scatter (from what? Electrons, Nuclear spins...) and produce... the Fourier Transform of the equal-time Correlation function. How?

$$\begin{aligned} |\tilde{\rho}(\mathbf{k})|^2 &= \tilde{\rho}(\mathbf{k})^* \tilde{\rho}(\mathbf{k}) = \int d\mathbf{x}' e^{i\mathbf{k}\cdot\mathbf{x}'} \rho(\mathbf{x}') \int d\mathbf{x} e^{-i\mathbf{k}\cdot\mathbf{x}} \rho(\mathbf{x}) \\ &= \int d\mathbf{x} d\mathbf{x}' e^{-i\mathbf{k}\cdot(\mathbf{x}-\mathbf{x}')} \rho(\mathbf{x}') \rho(\mathbf{x}) \\ &= \int d\mathbf{r} e^{-i\mathbf{k}\cdot\mathbf{r}} \int d\mathbf{x}' \rho(\mathbf{x}') \rho(\mathbf{x}' + \mathbf{r}) \\ &= \int d\mathbf{r} e^{-i\mathbf{k}\cdot\mathbf{r}} V \langle \rho(\mathbf{x}) \rho(\mathbf{x} + \mathbf{r}) \rangle = V \int d\mathbf{r} e^{-i\mathbf{k}\cdot\mathbf{r}} C(\mathbf{r}) \\ &= V \tilde{C}(\mathbf{k}). \end{aligned}$$




Fig. 10.6 X-ray scattering. A beam of wavevector \mathbf{k}_0 scatters off a density variation $\rho(\mathbf{x})$ with wavevector \mathbf{k} to a final wavevector $\mathbf{k}_0 + \mathbf{k}$; the intensity of the scattered beam is proportional to $|\tilde{\rho}(\mathbf{k})|^2$ [9, chapter 6].

Ideal gases: equal time correlations

Easiest, illustrative case (with no correlations). We need to compute from the FE free energy and density the fluctuations, and then consider what happens if we break the system into many sub-volumes (un-correlated).

Hemholtz' FE
and its derivative

$$\mathcal{F}^{\text{ideal}}(\rho(\mathbf{x}), T) = \rho(\mathbf{x}) k_B T [\log(\rho(\mathbf{x}) \lambda^3) - 1]$$

$$\alpha = \left. \frac{\partial^2 \mathcal{F}}{\partial \rho^2} \right|_{\rho_0} = k_B T / \rho_0 = P_0 / \rho_0^2$$

$$\mathcal{F}(\rho) = \frac{1}{2} \alpha (\rho - \rho_0)^2, \quad P\{\rho(\mathbf{x})\} \propto e^{-\beta \int \frac{1}{2} \alpha (\rho - \rho_0)^2 d\mathbf{x}}$$

Distributions of free energy and
density fluctuations in equilibrium

$$\langle (\rho - \rho_0)^2 \rangle = \frac{1}{\beta \alpha \Delta V}$$

Dirac' delta-function: no correlations.

$$C^{\text{ideal}}(\mathbf{r}, 0) = \frac{1}{\beta \alpha} \delta(\mathbf{r}).$$

Enter Onsager...

Lars Onsager

Lars Onsager (November 27, 1903
– October 5, 1976): Norwegian
physicist/chemist.

Known for: electrolytes... phase
transitions... Onsager relations....

Nobel prize (in Chemistry) in 1968.



Enter Onsager...

How to treat deviations from the equilibrium (read: correlations)?

O's regression hypothesis:

... we may assume that a spontaneous deviation from the equilibrium decays according to the same laws as one that has been produced artificially.

Average over initial conditions, thermal history. We get for the C the diffusion equation, again:

$$\begin{aligned} \frac{\partial C^{\text{ideal}}}{\partial t} &= \left\langle \frac{\partial}{\partial t} [\rho(\mathbf{r}, t)]_{\rho_i} (\rho_i(\mathbf{0}) - \rho_0) \right\rangle_{\text{eq}} \\ &= \left\langle D \nabla^2 [\rho(\mathbf{r}, t)]_{\rho_i} (\rho_i(\mathbf{0}) - \rho_0) \right\rangle_{\text{eq}} \\ &= D \nabla^2 \left\langle [\rho(\mathbf{r}, t)]_{\rho_i} (\rho_i(\mathbf{0}) - \rho_0) \right\rangle_{\text{eq}} \\ &= D \nabla^2 \langle (\rho(\mathbf{r}, t) - \rho_0)(\rho(\mathbf{0}, 0) - \rho_0) \rangle_{\text{ev}} \\ &= D \nabla^2 C^{\text{ideal}}(\mathbf{r}, t). \end{aligned}$$

$$\frac{\partial}{\partial t} [\rho(\mathbf{x}, t)]_{\rho_i} = D \nabla^2 [\rho(\mathbf{x}, t)]_{\rho_i}.$$

$$\begin{aligned} C(\mathbf{r}, \tau) &= \langle (\rho(\mathbf{x} + \mathbf{r}, t + \tau) - \rho_0)(\rho(\mathbf{x}, t) - \rho_0) \rangle_{\text{ev}} \\ &= \langle (\rho(\mathbf{r}, \tau) - \rho_0)(\rho(\mathbf{0}, 0) - \rho_0) \rangle_{\text{ev}} \\ &= \left\langle ([\rho(\mathbf{r}, \tau)]_{\rho_i} - \rho_0)(\rho_i(\mathbf{0}) - \rho_0) \right\rangle_{\text{eq}}. \end{aligned}$$

Example: 3D ideal gas from the DE

$$C^{\text{ideal}}(\mathbf{r}, \tau) = \frac{1}{\beta \alpha} G(\mathbf{r}, \tau) = \frac{1}{\beta \alpha} \left(\frac{1}{\sqrt{4\pi D \tau}} \right)^3 e^{-\mathbf{r}^2/4D|\tau|}.$$




Fig. 10.7 Noisy decay of a fluctuation. An unusual fluctuation at $t = 0$ will slowly decay to a more typical thermal configuration at a later time τ .




Fig. 10.8 Deterministic decay of an initial state. An initial condition with the same density will slowly decay to zero.

Susceptibility and linear response

The idea: define a measure for the response to a perturbation.

We assume that this can be measured “based on the past” via a response function χ . Note how and why this is linear (in f).

Then FT everything, and call χ as *the AC susceptibility* (language of magnets).

(Electricity: polarizability,

Magnetism: susceptibility again)

$$F_f(t) = - \int d\mathbf{x} f(\mathbf{x}, t) s(\mathbf{x}, t).$$

$$s(\mathbf{x}, t) = \int d\mathbf{x}' \int_{-\infty}^t dt' \chi(\mathbf{x} - \mathbf{x}', t - t') f(\mathbf{x}', t').$$

$$\tilde{s}(\mathbf{k}, \omega) = \tilde{\chi}(\mathbf{k}, \omega) \tilde{f}(\mathbf{k}, \omega),$$

Dissipation

χ splits into real and imaginary parts and $\text{Im } \chi$ relates to the “lag” of the response and to the dissipation per cycle (oscillatory force).

The zero-frequency limit (electrical analogue) relates the conductivity to limit of the polarizability.

$$\tilde{\chi}(\mathbf{k}, \omega) = \int d\mathbf{x} dt e^{i\omega t} e^{-i\mathbf{k}\cdot\mathbf{x}} \chi(\mathbf{x}, t) = \chi'(\mathbf{k}, \omega) + i\chi''(\mathbf{k}, \omega)$$

$$\begin{aligned} p(\omega) &= \frac{\omega |f_\omega|^2}{2} \int_{-\infty}^{\infty} d\tau \chi(\tau) \sin(\omega\tau) = \frac{\omega |f_\omega|^2}{2} \text{Im}[\tilde{\chi}(\omega)] \\ &= \frac{\omega |f_\omega|^2}{2} \chi''(\omega). \end{aligned}$$

$$\sigma = \lim_{\omega \rightarrow 0} \omega^2 \tilde{\alpha}(\omega)$$

Static susceptibility

Define via perturbed equilibrium
(no time-dependence).

Fluctuation-dissipation relation:
susceptibility vs. correlation
function in the zero frequency
limit.

Relation of these to fluctuations in
equilibrium and their (non-
extensive) scaling.

$$s(\mathbf{x}) = \int d\mathbf{x}' \chi_0(\mathbf{x} - \mathbf{x}') f(\mathbf{x}').$$

$$\chi_0(\mathbf{r}) = \beta C(\mathbf{r}, 0).$$

$$\tilde{\chi}_0(\mathbf{k}) = \tilde{\chi}(\mathbf{k}, \omega = 0).$$

$$\begin{aligned} k_B T \tilde{\chi}_0(\mathbf{k} = \mathbf{0}) &= \hat{C}(\mathbf{k} = \mathbf{0}, t = 0) = \int d\mathbf{r} \langle s(\mathbf{r} + \mathbf{x}) s(\mathbf{x}) \rangle \\ &= \int d\mathbf{r} \frac{1}{V} \left\langle \int d\mathbf{x} s(\mathbf{r} + \mathbf{x}) s(\mathbf{x}) \right\rangle \\ &= V \left\langle \frac{1}{V} \int d\mathbf{r}' s(\mathbf{r}') \frac{1}{V} \int d\mathbf{x} s(\mathbf{x}) \right\rangle \\ &= V \langle \langle s \rangle_{\text{space}}^2 \rangle. \end{aligned}$$

Fluctuation-dissipation theorem

Susceptibility χ relates to the correlations, thus the field and its fluctuations.

$$\chi(\mathbf{x}, t) = -\beta \frac{\partial C(\mathbf{x}, t)}{\partial t} \quad (t > 0).$$

In frequency domain, the imaginary part does the same.

$$\chi''(\omega) = \frac{\beta\omega}{2} \tilde{C}(\omega).$$

Thus also dissipated power: fluctuations are related to dissipation.

[FYI: there is a large universe of attempts to use this in out of equilibrium systems: measure χ and C, in order to define an effective temperature β_{eff} .]

$$\begin{aligned} p(\omega) &= \frac{\omega|f_\omega|^2}{2} \chi''(\omega) = \frac{\omega|f_\omega|^2}{2} \frac{\beta\omega}{2} \tilde{C}(\omega) \\ &= \frac{\beta\omega^2|f_\omega|^2}{4} \tilde{C}(\omega). \end{aligned}$$

Role of causality

The FT (frequency-dependent) susceptibility has real and imaginary parts: two functions instead of one ($\chi(t)$).

This can be used (Kramers-König -relation) to relate these to each other. The derivation follows from Cauchy's theorem in complex analysis (with the K-K contour).

$$\chi'(\omega) = \text{Re}[\tilde{\chi}(\omega)] = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\text{Im}[\tilde{\chi}(\omega')]}{\omega' - \omega} d\omega' = \frac{1}{\pi} \int_{-\infty}^{\infty} \frac{\chi''(\omega')}{\omega' - \omega} d\omega'.$$

$$\chi''(\omega) = -\frac{2\omega}{\pi} \int_0^{\infty} \chi'(\omega') \frac{1}{\omega'^2 - \omega^2} d\omega'.$$




Fig. 10.12 Kramers-Krönig contour. A contour C_ω in the complex ω' plane. The horizontal axis is $\text{Re}[\omega']$ and the vertical axis is $\text{Im}[\omega']$. The integration contour runs along the real axis from $-\infty$ to ∞ with a tiny semicircular detour near a pole at ω . The contour is closed with a semicircle back at infinity, where $\chi(\omega')$ vanishes rapidly. The contour encloses no singularities, so Cauchy's theorem tells us the integral around it is zero.

Homework

2.3 Generating random walks (Sethna 2.5 p. 28) HOMEWORK (5 points)

Please note that this exercise is computational, so in order to get help with possible problems, take a laptop to the exercise session or alternatively send your code and problem in advance to the TA. The preferred programming tool to use (from the point of view of debugging and getting TA help) is Python, but also others are acceptable.

(a) Write a routine to generate an N -step random walk in d dimensions, with each step uniformly distributed in the range $(-1/2, 1/2)$ in each dimension. (Generate the steps first as an $[N \times d]$ array, then do a cumulative sum.) Plot x_t versus t for a few 10 000-step random walks. Plot x versus y for a few two-dimensional random walks, with $N = 10, 1000, 100000$. (Try to keep the aspect ratio of the XY plot equal to one.) Does multiplying the number of steps by one hundred roughly increase the net distance by ten?

Each random walk is different and unpredictable, but the ensemble of random walks has elegant, predictable properties.

(b) Write a routine to calculate the endpoints of W random walks with N steps each in d dimensions. Do a scatter plot of the endpoints of 10000 random walks with $N = 1$ and 10, superimposed on the same plot. Notice that the longer random walks are distributed in a circularly symmetric pattern, even though the single step random walk $N = 1$ has a square probability distribution (arising from the single step range, see Fig 2.10 from Sethna p. 28).

This is an emergent symmetry; even though the walker steps longer distances along the diagonals of a square, a random walk several steps long has nearly perfect rotational symmetry. The most useful property of random walks is the central limit theorem. The endpoints of an ensemble of N step one-dimensional random walks with root-mean-square (RMS) step-size a has a Gaussian or normal probability distribution as $N \rightarrow \infty$,

$$\rho(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp(-x^2/2\sigma^2), \quad (1)$$

with $\sigma = \sqrt{Na}$.

(c) Calculate the RMS step-size a for one-dimensional steps uniformly distributed in $(-1/2, 1/2)$. Write a routine that plots a histogram of the endpoints of W one-dimensional random walks with N steps and 50 bins, along with the prediction of above equation for x in $(-3\sigma, 3\sigma)$. Do a histogram with $W = 10000$ and $N = 1, 2, 3, 5$. How quickly does the Gaussian distribution become a good approximation to the random walk?

Take home

This lecture looks at the classical measures of correlations and their decay. We shall get back to these topics later on, but you should read through the chapter and think of conditional probabilities. Read first the Chapter and check then the lecture slides again.

The take home consists of answering to the following three questions:

Give an example of X and Y that are correlated but there is no causal relation (X because of Y or X because of Y happened before) between them.

Take a (time) series of the binary kind $0110110011000111\dots$ (or subtract $-1/2$ from all the values so that the average might become zero). When would this be correlated?

Take instead a series like this: $\dots 0000111111(\dots)111000\dots$. This is clearly not a random one. Now start tossing a coin (0/1) and replace according to each toss one of the values with the new one. Does this correspond to the Onsager hypothesis and why? If the coin is biased, does the process relate to linear response?