

# MS-E2112 Multivariate Statistical Analysis (5cr)

## Lecture 8: Canonical Correlation Analysis

Lecturer: Pauliina Ilmonen  
Slides: Ilmonen/Kantala

# Contents

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

# Canonical Correlation Analysis

# Canonical Correlation Analysis

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Canonical correlation analysis involves partition of variables into two vectors  $x$  and  $y$ . The aim is to find linear combinations  $\alpha^T x$  and  $\beta^T y$  that have the largest possible correlation.

# Canonical Correlation Analysis

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Let  $x$  be a  $p$ -variate random vector and let  $y$  be a  $q$ -variate random vector. The object in canonical correlation analysis is to find linear combinations

$$u_k = \alpha_k^T x$$

and

$$v_k = \beta_k^T y$$

that maximizes the correlation  $|\text{corr}(u_k, v_k)|$  between  $u_k$  and  $v_k$  subject to

$$\text{var}(u_k) = \text{var}(v_k) = 1,$$

and

$$\text{corr}(u_k, u_t) = 0, \text{corr}(v_k, v_t) = 0, t < k.$$

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Quick reminder:

$$\text{corr}(w_1, w_2) = \frac{E[(w_1 - \mu_{w_1})(w_2 - \mu_{w_2})]}{\sigma_{w_1} \sigma_{w_2}}.$$

# Canonical Correlation Analysis

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

The vectors  $\alpha_k$  and  $\beta_k$  are called the  $k$ th canonical vectors and

$$\rho_k = |\text{corr}(u_k, v_k)|$$

are called canonical correlations.

# Canonical Correlation Analysis

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Whereas **principal component analysis** considers interrelationships **within a set of variables**, **canonical correlation analysis** considers relationships **between two groups of variables**.



# Canonical Correlation Analysis, Examples

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

- Exercise — health.
- Open book exams — closed book exams.
- Job satisfaction — performance.

# Canonical Correlation Analysis, Regression Analysis

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Canonical correlation analysis can be seen as an extension of multivariate regression analysis. However, note that in canonical correlation analysis there is **no assumption of causal asymmetry** -  $x$  and  $y$  are treated symmetrically!

# Canonical Correlation Analysis, Solution

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Let  $z = (x^T, y^T)^T$ , and let

$$\text{cov}(z) = \Sigma = \begin{bmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{bmatrix}.$$

Define

$$M_1 = \Sigma_{11}^{-1} \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21},$$

and

$$M_2 = \Sigma_{22}^{-1} \Sigma_{21} \Sigma_{11}^{-1} \Sigma_{12}.$$

# Canonical Correlation Analysis, Solution

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Now, the canonical vectors  $\alpha_k$  are the eigenvectors of  $M_1$  ( $\alpha_k$  corresponds to the  $k$ th largest eigenvalue), the canonical vectors  $\beta_k$  are the eigenvectors of  $M_2$ , and  $\rho_k^2$  are the eigenvalues of the matrix  $M_1$  (and of  $M_2$  as well). The proof of this solution can be found from pages 283-284 of [1].

# Canonical Correlation Analysis, Solution

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Note that the eigenvectors  $\alpha_k$  and  $\beta_k$  do not have length= 1!

Requirements

$$\text{var}(u_k) = \text{var}(\alpha_k^T x) = 1$$

and

$$\text{var}(v_k) = \text{var}(\beta_k^T y) = 1$$

define the lengths of the eigenvectors.

# Canonical Correlation Analysis, Solution

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

If the covariance matrices  $\Sigma_{11}$  and  $\Sigma_{22}$  are not full rank, similar results may be obtained using generalized inverses. One may also consider dimension reduction as a first step.

# Canonical Correlation Analysis, Sample Version

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Sample estimates  $\hat{\alpha}_k$ ,  $\hat{\beta}_k$  and  $\hat{\rho}_k$  of  $\alpha_k$ ,  $\beta_k$  and  $\rho_k$ , respectively, are obtained by using sample covariance matrices calculated from the samples  $x_1, x_2, \dots, x_n$ ,  $y_1, y_2, \dots, y_n$  and  $z_1, z_2, \dots, z_n$ .

# Canonical Correlation Analysis, Standardization

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

As in PCA, also in canonical correlation analysis, the data is sometimes standardized first.



## Testing Independence

# Testing Independence

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Assume that  $z = (x^T, y^T)^T \sim N_{p+q}(\mu, \Sigma)$ . Consider testing

$H_0$  :  $x$  and  $y$  are independent,

against

$H_1$  :  $x$  and  $y$  are not independent.

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

# Testing Independence

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Let  $m = \min\{p, q\}$ , and let

$$T = -\left(n - \frac{1}{2}(p + q + 3)\right) \ln\left(\prod_{k=1}^m (1 - \hat{\rho}_k^2)\right).$$

Now, under  $H_0$  (and under the assumption of multivariate normality), the test statistic  $T$  is asymptotically distributed as  $\chi^2(pq)$ .

# Testing Partial Independence

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Assume that  $z = (x^T, y^T)^T \sim N_{p+q}(\mu, \Sigma)$ . Consider testing

$H_0$  : Only  $s$  of the canonical correlation coefficients are nonzero,

against

$H_1$  : The number of nonzero canonical correlation coefficients is larger than  $s$ .

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

# Testing Partial Independence

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Let  $m = \min\{p, q\}$ , and let

$$T_s = -\left(n - \frac{1}{2}(p + q + 3)\right) \ln\left(\prod_{k=s+1}^m (1 - \hat{\rho}_k^2)\right).$$

Now, under  $H_0$  (and under the assumption of multivariate normality), the test statistic  $T$  is asymptotically distributed as  $\chi^2((p - s)(q - s))$ .

## Scoring and Predicting

# Scoring and Predicting

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Let  $X$  and  $Y$  denote the  $n \times p$  and  $n \times q$  data matrices for  $n$  individuals, and let  $\hat{\alpha}_k$  and  $\hat{\beta}_k$  denote the  $k$ th (sample) canonical vectors. Then the  $n \times 1$  vectors

$$\eta_k = X\hat{\alpha}_k$$

and

$$\phi_k = Y\hat{\beta}_k$$

denote the scores of the  $n$  individuals on the  $k$ th canonical correlation variables.

# Scoring and Predicting

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

If the  $x$  and  $y$  variables are interpreted as the "predictor" and "predicted" variables, respectively, then the  $\eta_k$  score vector can be used to predict the  $\phi_k$  score vector by using least square regression:

$$(\tilde{\phi}_k)_i = \hat{\rho}_k((\eta_k)_i - \hat{\alpha}_k^T \bar{x}) + \hat{\beta}_k^T \bar{y}.$$

The canonical correlation  $\hat{\rho}_k$  estimates the proportion of the variance of  $\phi_k$  that is explained by the regression on  $x$ .



# Example

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Example: closed book exams — open book exams.

# Example

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Marks in open-book (O) and closed-book (C) exams:

i	Mechanics (C)	Vectors (C)	Algebra (O)	Analysis (O)	Statistics (O)
1	77	82	67	67	81
2	63	78	80	70	81
3	75	73	71	66	81
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
100	46	52	53	41	40

Source: K. V. Mardia, J. T. Tent, J. M. Bibby, Multivariate analysis, Academic Press, London, 2003 (reprint of 1979).

# Example

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Means:

Variable	Mean
$x_1$	38.9545
$x_2$	50.5909
$y_1$	50.6023
$y_2$	46.6818
$y_3$	42.3068

# Example

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Covariance matrix

$$\Sigma =$$

$\Sigma_{11}$		$\Sigma_{12}$		
302.3	125.8	100.4	105.1	116.1
	170.9	84.2	93.6	97.9
		111.6	110.8	120.5
			217.9	153.8
				294.4
$\Sigma_{21}$		$\Sigma_{22}$		

# Example

Calculate the eigenvectors

$$M_1 = \Sigma_{11}^{-1} \Sigma_{12} \Sigma_{22}^{-1} \Sigma_{21} \Rightarrow \hat{\alpha}_k$$

and

$$M_2 = \Sigma_{22}^{-1} \Sigma_{21} \Sigma_{11}^{-1} \Sigma_{12} \Rightarrow \hat{\beta}_k.$$

Here

$$\hat{\alpha}_1 = \begin{bmatrix} 0.0260 \\ 0.0518 \end{bmatrix}$$

and

$$\hat{\beta}_1 = \begin{bmatrix} 0.0824 \\ 0.0081 \\ 0.0035 \end{bmatrix}.$$

# Example

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

$$u_1 = 0.0260x_1 + 0.0518x_2$$

and

$$v_1 = 0.0824y_1 + 0.0081y_2 + 0.0035y_3.$$

The highest correlation occurs between an average of  $x_1$  and  $x_2$  weighted on  $x_2$  and an average of  $y_1$ ,  $y_2$  and  $y_3$ , heavily weighted on  $y_1$

The canonical correlations

$$\rho_1 = 0.6630$$

and

$$\rho_2 = 0.0412.$$

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

# Example

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Predicting

$$\left(\tilde{\phi}_k\right)_i = \hat{\rho}_k \left( (\eta_k)_i - \hat{\alpha}_k^T \bar{x} \right) + \hat{\beta}_k^T \bar{y}.$$

Here

$$\begin{aligned} \left(\tilde{\phi}_1\right)_i &= 0.6630 \left( (\eta_1)_i - (0.0260 * 38.9545 + 0.0518 * 50.5909) \right) \\ &\quad + (0.0824 * 50.6023 + 0.0081 * 46.6818 + 0.0035 * 42.3068) \\ &\approx 0.6630(\eta_1)_i + 2.2905 \\ &\approx 0.6630(0.0260(x_1)_i + 0.0518(x_2)_i) + 2.2905 \\ &\approx 0.0172(x_1)_i + 0.0343(x_2)_i + 2.2905. \end{aligned}$$

Note that this almost predicts  $y_1$ .

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

## Words of Warning



# Some Words of Warning

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

- The procedure maximizes the correlation between the linear combination of variables — it can be more than difficult to interpret the results.
- Correlation does not automatically imply causality.
- Normality assumption is required in the tests presented in these slides.

# Next Week

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala

Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

Next week we will talk about discriminant analysis and classification.

## References

# References I

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala


Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

 K. V. Mardia, J. T. Kent, J. M. Bibby, *Multivariate Analysis*, Academic Press, London, 2003 (reprint of 1979).

# References II

Lecturer:  
Pauliina Ilmonen  
Slides:  
Ilmonen/Kantala


Canonical Correlation  
Analysis

Testing Independence

Scoring and Predicting

Words of Warning

References

-  R. V. Hogg, J. W. McKean, A. T. Craig, Introduction to Mathematical Statistics, Pearson Education, Upper Sadle River, 2005.
-  R. A. Horn, C. R. Johnson, Matrix Analysis, Cambridge University Press, New York, 1985.
-  R. A. Horn, C. R. Johnson, Topics in Matrix Analysis, Cambridge University Press, New York, 1991.