


General information

The exercise sessions will be held as blackboard sessions, where the participants will present their solutions to the group. As such, the problems should be set up and solved before the session. The focus of the exercises lies on analyzing and discussing the task at hand together with the group: thus, a perfect solution is not required to be awarded points. The (attempted) solutions should be submitted via email to the assistant at the start of the exercise session on February 10th. A point will be awarded for each question, and a person will be chosen to present their solution from the pool.

This exercise set concentrates on tokamaks and highlights especially the key concept of trapped particles that play a role in several physical processes in tokamaks, including transport of heat and particles.


Exercise 1.

Tritium breeding in tokamaks For tokamak reactors, a breeding blanket is expected to surround the plasma to breed tritium out of lithium through the following reactions:



Consider a 50-50 D-T fuel fusion plasma, neglecting the residual D-D fusion energy production. Estimate the contribution of the energy production by the n-Li reactions in the breeding blanket on the overall energy production in the reactor, assuming:

- a natural isotope ratio between ${}^6\text{Li}$ (7.42 %) and ${}^7\text{Li}$ (92.58 %).
- a 1:1 isotope ratio.
- Comment on your results.



vaikutusalat JENDL Dosimetry File 99, JAERI

Figure 1: Cross-sections for n-Li reactions.

Exercise 2.

Plasma confinement in tokamaks

In order to understand the plasma confinement in tokamaks, one must consider the loss- and transport phenomenon of heat and particles in the plasma. Classical and neoclassical predictions for radial transport in tokamaks will be investigated. These theoretical findings will be compared to experimentally observed values. Despite the many calculations and the platitude of information, the mathematics are straightforward, so do not lose hope.

- a) The experimentally observed confinement time in tokamaks is $\tau_E \sim 1$ s, while the minor radii in reactor scale tokamaks are approximately $a \sim 1$ m. Based on these ballpark values, estimate the radial transport (diffusion) coefficient for tokamaks, assuming $D \sim r^2/\tau$.
- b) Using the classical concept of collisional diffusion, the diffusion coefficient can be expressed as $D \sim (\text{step size})^2 \times (\text{collision frequency})$. Thus, the classical estimate for the radial diffusion coefficient is given by

$$D_{\text{classic}} \sim (r_{L,e})^2 \nu_{ei},$$

where $r_{L,e}$ is the electron gyroradius and ν_{ei} is the electron-ion collision frequency. The classical collisional thermal conductivity coefficients are

$$\chi_{s,\text{classic}} \sim (r_{L,s})^2 \nu_{ss}$$

where the subscript s refers to electrons and ions ($s \in \{i, e\}$), $r_{L,s}$ is the ion/electron gyro radius, while ν_{ss} are the ion-ion/electron-electron self-collision frequencies. Now, calculate the resulting values for D_{classic} , $\chi_{i,\text{classic}}$, and $\chi_{e,\text{classic}}$ assuming $\nu_{ei} \sim 4.6 \times 10^3 \text{ s}^{-1} \sim \nu_{ee}$, $\nu_{ii} \sim (m_e/m_i)^{1/2} \nu_{ei}$, $T_e = T_i \sim 5 \text{ keV}$, and $B_0 \sim 5 \text{ T}$.

- c) Due to the geometry of the tokamak, the toroidal field coils are denser towards the axis of symmetry, resulting in a stronger magnetic field closer to the axis of symmetry. Thus, the side of the plasma close to the axis of symmetry is referred to as the high field side (HFS), and the outer plasma as the low field side (LFS). This inhomogeneity in the toroidal magnetic field further enhances the transport in the device, due to cross-field drifts. These drifts effectively increase the diffusion step size, and the resulting transport, from the classical step size, and the resulting transport is referred to as neoclassical transport.

The radial gradient in the magnetic field also results in a phenomenon called particle trapping, in which a fraction of the particles are reflected before reaching the HFS plasma. Most of these trapped particles complete banana shaped orbits and are, naturally, called banana trapped particles, while the non-trapped particles are referred to as passing particles. Figure 2 should help, but watching an animation is even better, e.g. <https://www.youtube.com/watch?v=XUhNium3VEo>.

Calculate the diffusion coefficient estimates for both trapped and passing particles, using the neoclassical transport coefficient approximation for passing particles:

$$\begin{aligned} D_{\text{neoclassic}}^p &\sim 4q^2 D_{\text{classic}} \\ \chi_{s,\text{neoclassic}}^p &\sim q^2 \chi_{s,\text{classic}} \end{aligned}$$

where q is the safety factor, a quantity related to the windedness of the magnetic field. The neoclassical transport coefficient for trapped particles can then be estimated as

$$\begin{aligned} D_{\text{neoclassic}}^t &\sim 2.2q^2(R_0/r)^{3/2} D_{\text{classic}} \\ \chi_{e,\text{neoclassic}}^t &\sim 0.89q^2(R_0/r)^{3/2} \chi_{e,\text{classic}} \\ \chi_{i,\text{neoclassic}}^t &\sim 0.68q^2(R_0/r)^{3/2} \chi_{i,\text{classic}} \end{aligned}$$

Calculate the neoclassical estimates for the radial transport coefficients assuming $R_0 \sim 3$ m, $r \sim 1$ m, and $q \sim 3$.

- d) What conclusions can you make regarding classical and neoclassical collisional diffusion with regards the estimated cross-field transport calculated in a)?




Figure 2: Illustration of trapped particle orbits in a tokamak. The phenomenon is not limited just to fusion devices but occurs also in e.g. Earth's magnetosphere.

Exercise 3.

Trapped particles

Consider a particle starting at the outer mid-plane (LFS) of the poloidal cross-section of the torus with velocity $\mathbf{v}_0 = v_{\parallel,0}\mathbf{b} + \mathbf{v}_{\perp,0}$, where $\parallel$ stands for parallel and $\perp$ for perpendicular relative to the magnetic field direction $\mathbf{b}$. The kinetic energy of the particle can similarly be divided into perpendicular and parallel components. Ref. [1] might prove helpful in these calculations.

- Calculate the particle trapping condition for the pitch $\left(\frac{v_{\parallel,0}}{v_0} < \sqrt{1 - \frac{B_{\min}}{B_{\max}}}\right)$. Consider the extreme cases of the particle being located at the LFS (minimum B) and bouncing HFS (maximum B), and use the conservation of energy and magnetic moment $\left(\mu = \frac{mv_{\perp}^2}{2B}\right)$ and the fact that the parallel velocity is momentarily zero at the bounce.
- Show that this condition is equivalent to the one given in the lecture slides $\left(\frac{v_{\parallel,0}}{v_0} < \sqrt{2\frac{r}{R_0}}\right)$ when assuming $R_0 \gg r$ and $B(x) = \frac{B_0 R_0}{R_0 + x}$ with $-r \leq x \leq r$. B_0 is the magnetic field at the magnetic axis (in the middle of the cross section of the torus).

- c) Assuming Maxwellian velocity distribution, integrate over the trapped cone in the velocity space in spherical coordinates to obtain the fraction of trapped particles, i.e. calculate

$$f_t = \frac{1}{n} \int_{\theta_c}^{\pi-\theta_c} \sin\theta d\theta \int_0^{2\pi} d\phi \int_0^\infty F_M(v) v^2 dv,$$

where θ corresponds to $\cos\theta = \frac{v_{||,0}}{v_0}$ and θ_c to the boundary between trapped and passing particles. Utilising the calculations from b), the answer should be of the form $f_t = \sqrt{2r/R_0}$. What does this say about the balance between trapped and passing particles in a tokamak, i.e. where are the trapped particles most likely to dominate?

Exercise 4.

Maximum allowable current density in a tokamak

The helicity of the magnetic field in a tokamak is typically described with the safety factor, q , which is defined as the number of full toroidal loops that the magnetic field completes during a single poloidal revolution. The name stems from the fact that with certain q values, the plasma becomes susceptible to magnetohydrodynamic (MHD) instabilities, which may eventually lead to disruption and shut down of the reactor.

- a) Find the maximum allowable current density in the tokamak, such that the minimum edge safety factor, in this case $q > 1$, is still within operational boundaries imposed by plasma instabilities. Assume the toroidal magnetic field to be $B_T \sim 7$ T, and the major radius to be $R \sim 7$ m. Assume a circular symmetric poloidal cross-section, large aspect-ratio tokamak, such that the safety factor can be approximated by the equation:

$$q = \frac{rB_T}{RB_\theta},$$

where r is the local minor radius of the tokamak, B_T is the total toroidal field magnitude, R is the major radius of the tokamak, and B_θ is the total magnitude of the poloidal field. The poloidal field is induced by a current running through the plasma, and so depends on the current density.

[1] J. P. Freidberg, Plasma Physics and Fusion Energy, Cambridge University Press, 2007, p.484-485