



Aalto University  
School of Electrical  
Engineering

# Space Instrumentation

Magnetism and applications  
7.6.2020

Prof. Eija Tanskanen

*Aalto University  
School of Electrical Engineering  
Department of Radio Science and Engineering*



Aalto University  
School of Electrical  
Engineering

# Today

## Ground-based space weather measurements:

Magnetometers & all-sky cameras

Magnetic measurements

Geomagnetic activity indices


# Brief history of magnetic measurements

- First scientific magnetometer 1832 (Gauss)
- First scientific satellite 1958 (Explorer I)
- First magnetotail observations 1965
- Continuous ground-based observations since 1966 (Kyoto AL)
- Solar wind observations since 1966, continuous L1 observations since 1997 (ACE)



# Detecting (geo)magnetic activity (and currents)



Geomagnetic activity can be detected by all-sky cameras, magnetometers, radars, riometers, ionosondes, satellites ...




... or by an astronaut from a space shuttle



# Geomagnetic activity detected by magnetometers



# Analog magnetometer from 1960's



# Modern ground-based magnetometers




Flux gate magnetometer  
for scientific measurements




Helicopter magnetometer  
for magnetic surveys



# Spacecraft magnetometers




Eija Tanskanen, 2016



Ingo Richter, 2010

# All-sky cameras



# Observatories


Mawson, Antarctica



Nurmijärvi, Suomi



Addis Ababa, Ethiopia



Canberra, Australia



Apia, Western Samoa



L'Aquila, Italy



Greenland




Dalat, Vietnam




# Magnetometer chains

- IMAGE network
- CARISMA (earlier CANOPUS)
- 210 CHAIN
- Greenland chain
- MAGDAS
- Scandinavian SME (only historical data).




# Magnetometer networks

12 Kyoto AL observatories





29 IMAGE observatories





Courtesy of Häkkinen

# Conjugate magnetic measurements




# Geomagnetic storm signature & detection



Geomagnetic storms detected by magnetometers close to magnetic equator, not exactly at the equator due to the equatorial electrojets.


# Equatorial electrojet





# Geomagnetic storm index: Dst index

Formed as an envelope curve from the 12 equatorial magnetometers.




Geomagnetic storm starting on 15<sup>th</sup> July 2000  
- with a peak amplitude 300 nT  
- duration 5 days


# Geomagnetic activity at high-latitudes during storms

Pulsations


Storm-time substorms



# Extreme and moderate activity





Storm-time  
substorm



Non-storm  
substorm

# Magnetospheric substorm

- Magnetospheric substorms i.e. Birkeland's polar elementary storms, auroral substorms, etc.




# One substorm definition

“Magnetic substorm is a transient process, in which a significant amount of energy is carried from the solar wind into the auroral ionosphere and magnetosphere”.

*McPherron et al. 1979*




# “Typical” auroral substorm

- All substorms are different, there is not “a normal” substorm. Statistical properties can be computed, but they need to be understood as average properties and not a single such substorm does not need to exist.
- A typical substorm signature: a negative bay in north-south (X) component of the terrestrial magnetic field.




# Westward electrojet index AL/IL/SML/CL

- Following AL description IL index is formed based on IMAGE ground-based magnetic measurements in UT-sector 16-03 UT.



# Latitudinal variation of substorms


- Substorms were categorized in the latitude bins according to the station where the maximum deviation of the X component was recorded
- Latitudinal zones from north to south (geogr. coord.)
  - north of  $76^\circ$
  - $73^\circ - 76^\circ$
  - $69^\circ - 73^\circ$
  - $65^\circ - 69^\circ$
  - south of  $65^\circ$






# Auroral oval during a storm and a substorm

Storms





Substorms



# Storm-time substorms and non-storm substorms


Site of maximum  $dH/dt$  i.e. substorm onset location is dramatically more north for non-storm than storm-time substorms.



# Substorm morphology

Typical storm-time substorm is about twice as intense and carries about 2.5 times more energy into the ionosphere than a typical non-storm substorm.

(b) Typical stormtime substorm



(a) Typical isolated substorm

