



Aalto University
School of Science

Step-pulse excitaiton of ferromagnetic resonance

Sampo Hämäläinen


26.11.2012

Bachelors thesis seminar

Outline

1. What is ferromagnetic resonance?
2. Theoretical model for ferromagnetic resonance
3. How to excite ferromagnetic resonance?
4. Pulse inductive microwave magnetometry
 - Theoretical predictions and measurements

What is ferromagnetic resonance?




- Precession of magnetization at high frequency (~ 2 GHz+)
- Member of magnetic resonance family
 - Electron paramagnetic resonance
 - Nuclear magnetic resonance
- Very sensitive to magnetic environment

Why?



- FMR measures the dynamic response of magnetization
- Scans local magnetic environment
- Frequency and damping is essential
 - Limits magnetic switching speed
 - Spin torque switching

“Simple model” for precessional motion




- Larmor precession

$$\frac{dM}{dt} = \mu \times B = -\gamma\mu_0 M \times H$$

- Solution:
 - Circular motion around equilibrium direction H
- Unphysical: no damping mechanisms

Landau-Lifshitz equation of motion

$$\frac{\partial \vec{M}}{\partial t} = -\gamma \mu_0 \vec{M} \times \vec{H} - \frac{\lambda}{M_s^2} \vec{M} \times (\vec{M} \times \vec{H})$$




- Introduce damping term
 - Larmor precession + damping
- Important: $|\vec{M}(t)| = \text{constant}$
- Difficult to solve
- Unphysical in high damping region

Excitation of ferromagnetic resonance

- Multiple excitation methods:
 - Forced oscillator
 - Force oscillatory motion at constant frequency
 - Measure absorption vs. frequency
 - Free oscillation decay
 - Push magnetization out of equilibrium
 - Measure the decay process

Pulse inductive microwave magnetometry



- Free oscillatory decay
- Bias magnetization parallel to the signal line
- Apply fast rise time perpendicular magnetic field
- Measure perpendicular inducted component of magnetization


Solving Landau-Lifshitz equation: step pulse excitation...

- Bessel functions, trigonometry, 3D vector algebra, general spherical coordinate transformations...



... Solution!

$$\mathcal{E} \propto \frac{dM_y}{dt} = \omega_p M_s \cos(\omega_p t) \beta e^{-t/\tau}$$




- Exponentially decaying sinusoid as expected
- Angular frequency:

$$\omega_p^2 = \gamma^2 \mu_0^2 \left(\frac{\partial^2 U}{\partial \phi^2} \right)_{\phi=\phi_0}$$



Solution for uniaxial anisotropy

$$f_0^2 = \mu_0^2 \left(\frac{\gamma}{2\pi} \right)^2 M_s (H_k \cos 2(\phi - \alpha) + H_p \sin \phi + H_b \cos \phi)$$



- Resonance frequency depends
 - Anisotropy
 - Angle
 - Biasing field
- If saturation magnetization is known, gyroscopic splitting factor can be extracted

Experimental results



Conclusion

- Ferromagnetic resonance is important in the study of dynamic properties of magnetic thin films
- Solutions to Landau-Lifshitz agrees with experimental measurements
- Future: Measure more interesting samples
 - Stripe domains



Questions?

