

# Lecture 9: Gaussian and Particle Smoothers

Simo Särkkä

# Learning Outcomes

- 1 Summary of the Last Lecture
- 2 Extended and linearized smoothers
- 3 Gaussian RTS Smoothing
- 4 Particle Smoothing
- 5 Rao-Blackwellized Particle Smoothing
- 6 Summary and Demonstration

# Summary of the Last Lecture

- **Bayesian smoothing** is used for computing estimates of state trajectories **given the measurements on the whole trajectory**.
- **Rauch-Tung-Striebel (RTS) smoother** is the closed form smoother for **linear Gaussian** models.
- RTSS is **fixed-interval smoother**, there are also **fixed-point and fixed-lag** smoothers.

# Non-Linear Smoothing Problem

- **Non-linear Gaussian** state space model:

$$\mathbf{x}_k = \mathbf{f}(\mathbf{x}_{k-1}) + \mathbf{q}_{k-1}$$

$$\mathbf{y}_k = \mathbf{h}(\mathbf{x}_k) + \mathbf{r}_k,$$

- We want to compute **Gaussian approximations** to the smoothing distributions:

$$p(\mathbf{x}_k | \mathbf{y}_{1:T}) \approx N(\mathbf{x}_k | \mathbf{m}_k^S, \mathbf{P}_k^S).$$

# Extended Rauch-Tung-Striebel Smoother Derivation

- The **approximate joint distribution** of  $\mathbf{x}_k$  and  $\mathbf{x}_{k+1}$  is

$$p(\mathbf{x}_k, \mathbf{x}_{k+1} \mid \mathbf{y}_{1:k}) = \mathcal{N} \left( \begin{bmatrix} \mathbf{x}_k \\ \mathbf{x}_{k+1} \end{bmatrix} \mid \mathbf{m}_1, \mathbf{P}_1 \right),$$

where

$$\mathbf{m}_1 = \begin{pmatrix} \mathbf{m}_k \\ \mathbf{f}(\mathbf{m}_k) \end{pmatrix}$$
$$\mathbf{P}_1 = \begin{pmatrix} \mathbf{P}_k & \mathbf{P}_k \mathbf{F}_x^\top(\mathbf{m}_k) \\ \mathbf{F}_x(\mathbf{m}_k) \mathbf{P}_k & \mathbf{F}_x(\mathbf{m}_k) \mathbf{P}_k \mathbf{F}_x^\top(\mathbf{m}_k) + \mathbf{Q}_k \end{pmatrix}.$$

- The rest of the derivation is **analogous to the linear RTS smoother**.

## Extended Rauch-Tung-Striebel Smoother

The equations for the extended RTS smoother are

$$\mathbf{m}_{k+1}^- = \mathbf{f}(\mathbf{m}_k)$$

$$\mathbf{P}_{k+1}^- = \mathbf{F}_x(\mathbf{m}_k) \mathbf{P}_k \mathbf{F}_x^\top(\mathbf{m}_k) + \mathbf{Q}_k$$

$$\mathbf{G}_k = \mathbf{P}_k \mathbf{F}_x^\top(\mathbf{m}_k) [\mathbf{P}_{k+1}^-]^{-1}$$

$$\mathbf{m}_k^s = \mathbf{m}_k + \mathbf{G}_k [\mathbf{m}_{k+1}^s - \mathbf{m}_{k+1}^-]$$

$$\mathbf{P}_k^s = \mathbf{P}_k + \mathbf{G}_k [\mathbf{P}_{k+1}^s - \mathbf{P}_{k+1}^-] \mathbf{G}_k^\top,$$

where the matrix  $\mathbf{F}_x(\mathbf{m}_k)$  is the Jacobian matrix of  $\mathbf{f}(\mathbf{x})$  evaluated at  $\mathbf{m}_k$ .

# Statistically Linearized Rauch-Tung-Striebel Smoother Derivation

- With **statistical linearization** we get the approximation

$$p(\mathbf{x}_k, \mathbf{x}_{k+1} | \mathbf{y}_{1:k}) = N \left( \begin{bmatrix} \mathbf{x}_k \\ \mathbf{x}_{k+1} \end{bmatrix} \mid \mathbf{m}_1, \mathbf{P}_1 \right),$$

where

$$\mathbf{m}_1 = \begin{pmatrix} \mathbf{m}_k \\ E[\mathbf{f}(\mathbf{x}_k)] \end{pmatrix}$$

$$\mathbf{P}_1 = \begin{pmatrix} \mathbf{P}_k & E[\mathbf{f}(\mathbf{x}_k) \delta \mathbf{x}_k^T]^T \\ E[\mathbf{f}(\mathbf{x}_k) \delta \mathbf{x}_k^T] & E[\mathbf{f}(\mathbf{x}_k) \delta \mathbf{x}_k^T] \mathbf{P}_k^{-1} E[\mathbf{f}(\mathbf{x}_k) \delta \mathbf{x}_k^T]^T + \mathbf{Q}_k \end{pmatrix}.$$

- The **expectations** are taken with respect to **filtering distribution** of  $\mathbf{x}_k$ .
- The derivation proceeds as with **linear RTS smoother**.

## Statistically Linearized Rauch-Tung-Striebel Smoother

The equations for the statistically linearized RTS smoother are

$$\mathbf{m}_{k+1}^- = E[\mathbf{f}(\mathbf{x}_k)]$$

$$\mathbf{P}_{k+1}^- = E[\mathbf{f}(\mathbf{x}_k) \delta \mathbf{x}_k^\top] \mathbf{P}_k^{-1} E[\mathbf{f}(\mathbf{x}_k) \delta \mathbf{x}_k^\top]^\top + \mathbf{Q}_k$$

$$\mathbf{G}_k = E[\mathbf{f}(\mathbf{x}_k) \delta \mathbf{x}_k^\top]^\top [\mathbf{P}_{k+1}^-]^{-1}$$

$$\mathbf{m}_k^s = \mathbf{m}_k + \mathbf{G}_k [\mathbf{m}_{k+1}^s - \mathbf{m}_{k+1}^-]$$

$$\mathbf{P}_k^s = \mathbf{P}_k + \mathbf{G}_k [\mathbf{P}_{k+1}^s - \mathbf{P}_{k+1}^-] \mathbf{G}_k^\top,$$

where the expectations are taken with respect to the filtering distribution  $\mathbf{x}_k \sim N(\mathbf{m}_k, \mathbf{P}_k)$ .



# Gaussian Rauch-Tung-Striebel Smoother Derivation

- With **Gaussian moment matching** we get the approximation

$$p(\mathbf{x}_k, \mathbf{x}_{k+1} | \mathbf{y}_{1:k}) = \mathcal{N} \left( \begin{bmatrix} \mathbf{x}_k \\ \mathbf{x}_{k+1} \end{bmatrix} \middle| \begin{bmatrix} \mathbf{m}_k \\ \mathbf{m}_{k+1}^- \end{bmatrix}, \begin{bmatrix} \mathbf{P}_k & \mathbf{D}_{k+1} \\ \mathbf{D}_{k+1}^\top & \mathbf{P}_{k+1}^- \end{bmatrix} \right),$$

where

$$\mathbf{m}_{k+1}^- = \int \mathbf{f}(\mathbf{x}_k) \mathcal{N}(\mathbf{x}_k | \mathbf{m}_k, \mathbf{P}_k) d\mathbf{x}_k$$

$$\mathbf{P}_{k+1}^- = \int [\mathbf{f}(\mathbf{x}_k) - \mathbf{m}_{k+1}^-] [\mathbf{f}(\mathbf{x}_k) - \mathbf{m}_{k+1}^-]^\top \\ \times \mathcal{N}(\mathbf{x}_k | \mathbf{m}_k, \mathbf{P}_k) d\mathbf{x}_k + \mathbf{Q}_k$$

$$\mathbf{D}_{k+1} = \int [\mathbf{x}_k - \mathbf{m}_k] [\mathbf{f}(\mathbf{x}_k) - \mathbf{m}_{k+1}^-]^\top \mathcal{N}(\mathbf{x}_k | \mathbf{m}_k, \mathbf{P}_k) d\mathbf{x}_k.$$

## Gaussian Rauch-Tung-Striebel Smoother

The equations for the Gaussian RTS smoother are

$$\mathbf{m}_{k+1}^- = \int \mathbf{f}(\mathbf{x}_k) \mathcal{N}(\mathbf{x}_k | \mathbf{m}_k, \mathbf{P}_k) d\mathbf{x}_k$$

$$\mathbf{P}_{k+1}^- = \int [\mathbf{f}(\mathbf{x}_k) - \mathbf{m}_{k+1}^-] [\mathbf{f}(\mathbf{x}_k) - \mathbf{m}_{k+1}^-]^\top \\ \times \mathcal{N}(\mathbf{x}_k | \mathbf{m}_k, \mathbf{P}_k) d\mathbf{x}_k + \mathbf{Q}_k$$

$$\mathbf{D}_{k+1} = \int [\mathbf{x}_k - \mathbf{m}_k] [\mathbf{f}(\mathbf{x}_k) - \mathbf{m}_{k+1}^-]^\top \mathcal{N}(\mathbf{x}_k | \mathbf{m}_k, \mathbf{P}_k) d\mathbf{x}_k$$

$$\mathbf{G}_k = \mathbf{D}_{k+1} [\mathbf{P}_{k+1}^-]^{-1}$$

$$\mathbf{m}_k^s = \mathbf{m}_k + \mathbf{G}_k (\mathbf{m}_{k+1}^s - \mathbf{m}_{k+1}^-)$$

$$\mathbf{P}_k^s = \mathbf{P}_k + \mathbf{G}_k (\mathbf{P}_{k+1}^s - \mathbf{P}_{k+1}^-) \mathbf{G}_k^\top.$$

- Recall the 3rd order spherical Gaussian integral rule:

$$\int \mathbf{g}(\mathbf{x}) N(\mathbf{x} | \mathbf{m}, \mathbf{P}) d\mathbf{x} \\ \approx \frac{1}{2n} \sum_{i=1}^{2n} \mathbf{g}(\mathbf{m} + \sqrt{\mathbf{P}} \boldsymbol{\xi}^{(i)}),$$

where

$$\boldsymbol{\xi}^{(i)} = \begin{cases} \sqrt{n} \mathbf{e}_i & , \quad i = 1, \dots, n \\ -\sqrt{n} \mathbf{e}_{i-n} & , \quad i = n+1, \dots, 2n, \end{cases}$$

where  $\mathbf{e}_i$  denotes a unit vector to the direction of coordinate axis  $i$ .

# Cubature Smoother Derivation [2/2]

- We get the approximation

$$p(\mathbf{x}_k, \mathbf{x}_{k+1} \mid \mathbf{y}_{1:k}) = N \left( \begin{bmatrix} \mathbf{x}_k \\ \mathbf{x}_{k+1} \end{bmatrix} \mid \begin{bmatrix} \mathbf{m}_k \\ \mathbf{m}_{k+1}^- \end{bmatrix}, \begin{bmatrix} \mathbf{P}_k & \mathbf{D}_{k+1} \\ \mathbf{D}_{k+1}^\top & \mathbf{P}_{k+1}^- \end{bmatrix} \right),$$

where

$$\mathcal{X}_k^{(i)} = \mathbf{m}_k + \sqrt{\mathbf{P}_k} \boldsymbol{\xi}^{(i)}$$

$$\mathbf{m}_{k+1}^- = \frac{1}{2n} \sum_{i=1}^{2n} \mathbf{f}(\mathcal{X}_k^{(i)})$$

$$\mathbf{P}_{k+1}^- = \frac{1}{2n} \sum_{i=1}^{2n} [\mathbf{f}(\mathcal{X}_k^{(i)}) - \mathbf{m}_{k+1}^-] [\mathbf{f}(\mathcal{X}_k^{(i)}) - \mathbf{m}_{k+1}^-]^\top + \mathbf{Q}_k$$

$$\mathbf{D}_{k+1} = \frac{1}{2n} \sum_{i=1}^{2n} [\mathcal{X}_k^{(i)} - \mathbf{m}_k] [\mathbf{f}(\mathcal{X}_k^{(i)}) - \mathbf{m}_{k+1}^-]^\top.$$

## Cubature Rauch-Tung-Striebel Smoother

- 1 Form the sigma points:

$$\mathcal{X}_k^{(i)} = \mathbf{m}_k + \sqrt{\mathbf{P}_k} \boldsymbol{\xi}^{(i)}, \quad i = 1, \dots, 2n,$$

where the unit sigma points are defined as

$$\boldsymbol{\xi}^{(i)} = \begin{cases} \sqrt{n} \mathbf{e}_i & , \quad i = 1, \dots, n \\ -\sqrt{n} \mathbf{e}_{i-n} & , \quad i = n+1, \dots, 2n. \end{cases}$$

- 2 Propagate the sigma points through the dynamic model:

$$\hat{\mathcal{X}}_{k+1}^{(i)} = \mathbf{f}(\mathcal{X}_k^{(i)}), \quad i = 1, \dots, 2n.$$

## Cubature Rauch-Tung-Striebel Smoother (cont.)

- ③ Compute the predicted mean  $\mathbf{m}_{k+1}^-$ , the predicted covariance  $\mathbf{P}_{k+1}^-$  and the cross-covariance  $\mathbf{D}_{k+1}$ :

$$\mathbf{m}_{k+1}^- = \frac{1}{2n} \sum_{i=1}^{2n} \hat{\mathcal{X}}_{k+1}^{(i)}$$

$$\mathbf{P}_{k+1}^- = \frac{1}{2n} \sum_{i=1}^{2n} (\hat{\mathcal{X}}_{k+1}^{(i)} - \mathbf{m}_{k+1}^-) (\hat{\mathcal{X}}_{k+1}^{(i)} - \mathbf{m}_{k+1}^-)^\top + \mathbf{Q}_k$$

$$\mathbf{D}_{k+1} = \frac{1}{2n} \sum_{i=1}^{2n} (\mathcal{X}_k^{(i)} - \mathbf{m}_k) (\hat{\mathcal{X}}_{k+1}^{(i)} - \mathbf{m}_{k+1}^-)^\top.$$

## Cubature Rauch-Tung-Striebel Smoother (cont.)

- 4 Compute the gain  $\mathbf{G}_k$ , mean  $\mathbf{m}_k^S$  and covariance  $\mathbf{P}_k^S$  as follows:

$$\mathbf{G}_k = \mathbf{D}_{k+1} [\mathbf{P}_{k+1}^-]^{-1}$$

$$\mathbf{m}_k^S = \mathbf{m}_k + \mathbf{G}_k (\mathbf{m}_{k+1}^S - \mathbf{m}_{k+1}^-)$$

$$\mathbf{P}_k^S = \mathbf{P}_k + \mathbf{G}_k (\mathbf{P}_{k+1}^S - \mathbf{P}_{k+1}^-) \mathbf{G}_k^T.$$

## Unscented Rauch-Tung-Striebel Smoother

- 1 Form the sigma points:

$$\begin{aligned}\mathcal{X}_k^{(0)} &= \mathbf{m}_k, \\ \mathcal{X}_k^{(i)} &= \mathbf{m}_k + \sqrt{n + \lambda} \left[ \sqrt{\mathbf{P}_k} \right]_i \\ \mathcal{X}_k^{(i+n)} &= \mathbf{m}_k - \sqrt{n + \lambda} \left[ \sqrt{\mathbf{P}_k} \right]_i, \quad i = 1, \dots, n.\end{aligned}$$

- 2 Propagate the sigma points through the dynamic model:

$$\hat{\mathcal{X}}_{k+1}^{(i)} = \mathbf{f}(\mathcal{X}_k^{(i)}), \quad i = 0, \dots, 2n.$$



## Unscented Rauch-Tung-Striebel Smoother (cont.)

- 3 Compute predicted mean, covariance and cross-covariance:

$$\mathbf{m}_{k+1}^- = \sum_{i=0}^{2n} W_i^{(m)} \hat{\mathcal{X}}_{k+1}^{(i)}$$

$$\mathbf{P}_{k+1}^- = \sum_{i=0}^{2n} W_i^{(c)} (\hat{\mathcal{X}}_{k+1}^{(i)} - \mathbf{m}_{k+1}^-) (\hat{\mathcal{X}}_{k+1}^{(i)} - \mathbf{m}_{k+1}^-)^\top + \mathbf{Q}_k$$

$$\mathbf{D}_{k+1} = \sum_{i=0}^{2n} W_i^{(c)} (\mathcal{X}_k^{(i)} - \mathbf{m}_k) (\hat{\mathcal{X}}_{k+1}^{(i)} - \mathbf{m}_{k+1}^-)^\top,$$

## Unscented Rauch-Tung-Striebel Smoother (cont.)

- 4 Compute gain smoothed mean and smoothed covariance:  
as follows:

$$\mathbf{G}_k = \mathbf{D}_{k+1} [\mathbf{P}_{k+1}^-]^{-1}$$

$$\mathbf{m}_k^S = \mathbf{m}_k + \mathbf{G}_k (\mathbf{m}_{k+1}^S - \mathbf{m}_{k+1}^-)$$

$$\mathbf{P}_k^S = \mathbf{P}_k + \mathbf{G}_k (\mathbf{P}_{k+1}^S - \mathbf{P}_{k+1}^-) \mathbf{G}_k^T.$$

# Other Gaussian RTS Smoothers

- **Gauss-Hermite RTS smoother** is based on multidimensional Gauss-Hermite integration.
- **Bayes-Hermite or Gaussian Process RTS smoother** uses Gaussian process based quadrature (Bayes-Hermite).
- **Monte Carlo integration based RTS smoothers.**
- **Central differences etc.**

# Particle Smoothing: Direct SIR

- The smoothing solution can be obtained from SIR by **storing the whole state histories** into the particles.
- **Special care** is needed on the **resampling** step.
- The **smoothed distribution approximation** is then of the form

$$p(\mathbf{x}_k | \mathbf{y}_{1:T}) \approx \sum_{i=1}^N w_T^{(i)} \delta(\mathbf{x}_k - \mathbf{x}_k^{(i)}),$$

where  $\mathbf{x}_k^{(i)}$  is the  $k$ th component in  $\mathbf{x}_{1:T}^{(i)}$ .

- Unfortunately, the approximation is often quite **degenerate**.

# Particle Smoothing: Backward Simulation [1/2]

- In **backward-simulation particle smoother** we simulate individual **trajectories backwards**.
- The simulated samples are drawn from the **particle filter samples**.
- Uses the previous filtering results in smoothing  $\Rightarrow$  **less degenerate** than the direct SIR smoother.
- **Idea:**
  - Assume now that we have already simulated  $\tilde{\mathbf{x}}_{k+1:T}$  from the smoothing distribution.
  - From the Bayesian smoothing equations we get

$$p(\mathbf{x}_k | \tilde{\mathbf{x}}_{k+1}, \mathbf{y}_{1:T}) \propto p(\tilde{\mathbf{x}}_{k+1} | \mathbf{x}_k) p(\mathbf{x}_k | \mathbf{y}_{1:k}).$$

## Backward simulation particle smoother

Given the weighted set of particles  $\{w_k^{(i)}, \mathbf{x}_k^{(i)}\}$  representing the filtering distributions:

- Choose  $\tilde{\mathbf{x}}_T = \mathbf{x}_T^{(i)}$  with probability  $w_T^{(i)}$ .
- For  $k = T - 1, \dots, 0$ :
  - 1 Compute new weights by

$$w_{k|k+1}^{(i)} \propto w_k^{(i)} p(\tilde{\mathbf{x}}_{k+1} | \mathbf{x}_k^{(i)})$$

- 2 Choose  $\tilde{\mathbf{x}}_k = \mathbf{x}_k^{(i)}$  with probability  $w_{k|k+1}^{(i)}$

Given  $S$  iterations resulting in  $\tilde{\mathbf{x}}_{1:T}^{(j)}$  for  $j = 1, \dots, S$  the smoothing distribution approximation is

$$p(\mathbf{x}_{1:T} | \mathbf{y}_{1:T}) \approx \frac{1}{S} \sum_j \delta(\mathbf{x}_{1:T} - \tilde{\mathbf{x}}_{1:T}^{(j)}).$$

# Particle Smoothing: Reweighting [1/2]

- The **reweighting particle smoother** is based on computing new weights  $w_{k+1|T}^{(i)}$  for the SIR filter particles such that:

$$p(\mathbf{x}_{k+1} | \mathbf{y}_{1:T}) \approx \sum_i w_{k+1|T}^{(i)} \delta(\mathbf{x}_{k+1} - \mathbf{x}_{k+1}^{(i)}).$$

- Recall the smoothing equation

$$p(\mathbf{x}_k | \mathbf{y}_{1:T}) = p(\mathbf{x}_k | \mathbf{y}_{1:k}) \int \left[ \frac{p(\mathbf{x}_{k+1} | \mathbf{x}_k) p(\mathbf{x}_{k+1} | \mathbf{y}_{1:T})}{p(\mathbf{x}_{k+1} | \mathbf{y}_{1:k})} \right] d\mathbf{x}_{k+1}$$

- We use SIR filter samples to form approximations as follows:

$$\int \frac{p(\mathbf{x}_{k+1} | \mathbf{x}_k) p(\mathbf{x}_{k+1} | \mathbf{y}_{1:T})}{p(\mathbf{x}_{k+1} | \mathbf{y}_{1:k})} d\mathbf{x}_{k+1} \approx \sum_i w_{k+1|T}^{(i)} \frac{p(\mathbf{x}_{k+1}^{(i)} | \mathbf{x}_k)}{p(\mathbf{x}_{k+1}^{(i)} | \mathbf{y}_{1:k})}$$
$$p(\mathbf{x}_{k+1} | \mathbf{y}_{1:k}) \approx \sum_j w_k^{(j)} p(\mathbf{x}_{k+1} | \mathbf{x}_k^{(j)})$$

## Reweighting Particle Smoother

Given the weighted set of particles  $\{w_k^{(i)}, \mathbf{x}_k^{(i)}\}$  representing the filtering distribution, we can form approximations to the marginal smoothing distributions as follows:

- Start by setting  $w_{T|T}^{(i)} = w_T^{(i)}$  for  $i = 1, \dots, n$ .
- For each  $k = T - 1, \dots, 0$  do the following:
  - Compute new importance weights by

$$w_{k|T}^{(i)} \propto \sum_j w_{k+1|T}^{(j)} \frac{w_k^{(i)} p(\mathbf{x}_{k+1}^{(i)} | \mathbf{x}_k^{(i)})}{\left[ \sum_l w_k^{(l)} p(\mathbf{x}_{k+1}^{(l)} | \mathbf{x}_k^{(l)}) \right]}.$$

At each step  $k$  the marginal smoothing distribution can be approximated as

$$p(\mathbf{x}_k | \mathbf{y}_{1:T}) \approx \sum_j w_{k|T}^{(j)} \delta(\mathbf{x}_k - \mathbf{x}_k^{(j)}).$$



# Rao-Blackwellized Particle Smoothing: Direct SIR

- Recall e.g. the hierarchical the **Rao-Blackwellized particle filtering model**:

$$\mathbf{u}_k \sim p(\mathbf{u}_k | \mathbf{u}_{k-1})$$

$$\mathbf{x}_k = \mathbf{A}(\mathbf{u}_{k-1}) \mathbf{x}_{k-1} + \mathbf{q}_k, \quad \mathbf{q}_k \sim N(\mathbf{0}, \mathbf{Q})$$

$$\mathbf{y}_k = \mathbf{H}(\mathbf{u}_k) \mathbf{x}_k + \mathbf{r}_k, \quad \mathbf{r}_k \sim N(\mathbf{0}, \mathbf{R})$$

- The direct SIR based **Rao-Blackwellized particle smoother**:
  - During filtering store the whole sampled **state and Kalman filter histories** to the particles.
  - At the smoothing step, apply **Rauch-Tung-Striebel smoothers** to each of the Kalman filter histories.
- The **smoothing distribution approximation**:

$$p(\mathbf{x}_k, \mathbf{u}_k | \mathbf{y}_{1:T}) \approx \sum_{i=1}^N w_T^{(i)} \delta(\mathbf{u}_k - \mathbf{u}_k^{(i)}) N(\mathbf{x}_k | \mathbf{m}_k^{s,(i)}, \mathbf{P}_k^{s,(i)}).$$

- Also has the **degeneracy** problem.

- The **RB backward-sampling smoother** can be implemented in many ways:
  - **Sample both the components** backwards (leads to a pure sample representation).
  - **Sample the latent variables only** – requires quite complicated backward Kalman filtering computations.
  - **Kim's approximation**: just use the plain backward-sampling to the latent variable marginal.
- The **RB reweighting particle smoothing** is not possible exactly, but can be approximated using the above ideas.

- **Extended, statistically linearized and unscented RTS smoothers** are the approximate nonlinear smoothers corresponding to EKF, SLF and UKF.
- **Gaussian RTS smoothers**: cubature RTS smoother, Gauss-Hermite RTS smoothers and various others
- **Particle smoothing** can be done by storing the whole **state histories** in SIR algorithm.
- **Rao-Blackwellized particle smoother** is a combination of particle smoothing and RTS smoothing.

- Pendulum model:

$$\begin{pmatrix} x_k^1 \\ x_k^2 \end{pmatrix} = \underbrace{\begin{pmatrix} x_{k-1}^1 + x_{k-1}^2 \Delta t \\ x_{k-1}^2 - g \sin(x_{k-1}^1) \Delta t \end{pmatrix}}_{\mathbf{f}(\mathbf{x}_{k-1})} + \begin{pmatrix} 0 \\ q_{k-1} \end{pmatrix}$$

$$y_k = \underbrace{\sin(x_k^1)}_{\mathbf{h}(\mathbf{x}_k)} + r_k,$$

- The required Jacobian matrix for ERTSS:

$$\mathbf{F}_x(\mathbf{x}) = \begin{pmatrix} 1 & \Delta t \\ -g \cos(x^1) \Delta t & 1 \end{pmatrix}$$

- The required expected value for SLRTSS is

$$E[\mathbf{f}(\mathbf{x})] = \begin{pmatrix} m_1 + m_2 \Delta t \\ m_2 - g \sin(m_1) \exp(-P_{11}/2) \Delta t \end{pmatrix}$$

- And the cross term:

$$E[\mathbf{f}(\mathbf{x}) (\mathbf{x} - \mathbf{m})^\top] = \begin{pmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{pmatrix},$$

where

$$c_{11} = P_{11} + \Delta t P_{12}$$

$$c_{12} = P_{12} + \Delta t P_{22}$$

$$c_{21} = P_{12} - g \Delta t \cos(m_1) P_{11} \exp(-P_{11}/2)$$

$$c_{22} = P_{22} - g \Delta t \cos(m_1) P_{12} \exp(-P_{11}/2)$$