

Aalto university

Björn Ivarsson

**Exercise sheet 7**

Complex Analysis, MS-C1300.

**Hand in exercise 1 and 2 for grading. Deadline Monday 16.11**

**at 23:59.** The exercises should be uploaded to the correct folder on MyCourses as one pdf-file with name and student number in the file name. **Submission via MyCourses is the only accepted way.**

Done during class Tuesday 17.11 or Wednesday 18.11.

- (1) By means of the local Cauchy Theorem (Cauchy's Theorem in a disk), assuming that the circles are positively oriented, calculate:

(a)

$$\int_{|z|=1} \sqrt{9-z^2} dz$$

(3p)

(b)

$$\int_{|z|=1} \frac{1}{z^2+2z} dz$$

(3p)

- (2) Let  $\gamma(t) = 2 \cos t + i \sin t$  for  $0 \leq t \leq 2\pi$ . Evaluate:

(a)

$$\int_{\gamma} \frac{1}{z} dz$$

(3p)

(b)

$$\int_{\gamma} \frac{1}{z^2+2iz} dz$$

(3p)

(*Hint:* Winding numbers and partial fractions help here.)

- (3) Calculate

$$\int_{|z+i|=3/2} \frac{1}{z^4+z^2} dz$$

where the trajectory  $|z+i|=3/2$  is positively oriented.

- (4) Let  $\gamma$  and  $\beta$  be closed, piecewise smooth paths in  $\mathbb{C}$  with the same initial point. Show that  $n(-\gamma, z) = -n(\gamma, z)$  for every

$z \in \mathbb{C} \setminus |\gamma|$  and that  $n(\gamma + \beta, z) = n(\gamma, z) + n(\beta, z)$  for every  
 $z \in \mathbb{C} \setminus (|\gamma| \cup |\beta|)$ .