


# PHYS-C0252 - Quantum Mechanics Part 5


16.11.2020-08.12.2020

Tapio.Ala-Nissila@aalto.fi



## 4.3 Finite Potential Barrier

- The first nontrivial case is that of a finite potential barrier, where the QM particle can penetrate in and scatter from:



The full solution of this problem would entail dynamical treatment, and according to the SE

$$\Psi(x, t) = \int c(E) \psi_E(x) e^{-iEt/\hbar}$$

To the left of the barrier

$$\psi''_E(x) = -k^2 \psi_E(x), \quad E = \frac{\hbar^2 k^2}{2m}$$

and the wave solution is

$$\psi_E(x) = A_I e^{ikx} + A_R e^{-ikx}$$

where the intensity of the *incident* wave is  $|A_I|^2$  and that of the *reflected* wave  $|A_R|^2$

- When the energy of the incoming particle is larger than that of the barrier (classical crossing)

$$\psi''_E(x) = -k_B^2 \psi_E(x), \quad E = \frac{\hbar^2 k_B^2}{2m} + V_B$$

whose general solution is

$$\psi_E(x) = Ae^{ik_B x} + A'e^{-ik_B x}$$

For  $E < V$ , the region is classically forbidden (reflection), but the SE gives

$$\psi_E''(x) = \beta^2 \psi_E(x), \quad E = -\frac{\hbar^2 \beta^2}{2m} + V_B$$

and the general solution becomes a decaying one

$$\psi_E(x) = Be^{-\beta x} + B'e^{\beta x}$$

Finally, on the r.h.s. of the barrier (equals l.h.s.)

$$\psi_E(x) = A_T e^{ikx}, \quad k = \sqrt{2Em}/\hbar$$

The physically interesting quantities here are the ratios of the reflected and transmitted intensities

$$R = \frac{|A_R|^2}{|A_I|^2} \quad \text{and} \quad T = \frac{|A_T|^2}{|A_I|^2}.$$

These are called *reflection* and *transmission probabilities* and  $R + T = 1$

We focus here on a particle whose energy is below the barrier:

$$\psi_E(x) = \begin{cases} A_I e^{+ikx} + A_R e^{-ikx} & \text{if } -\infty < x < 0 \\ B e^{-\beta x} + B' e^{+\beta x} & \text{if } 0 < x < a \\ A_T e^{+ikx} & \text{if } a < x < \infty, \end{cases}$$

Continuity at  $x = 0$  and  $a$  gives

$$A_I + A_R = B + B' \quad \text{and} \quad ikA_I - ikA_R = -\beta B + \beta B',$$

$$B e^{-\beta a} + B' e^{+\beta a} = A_T e^{ika} \quad \text{and} \quad -\beta B e^{-\beta a} + \beta B' e^{+\beta a} = ikA_T e^{ika}$$

from which we can get the amplitudes as a function of  $B$ :

$$2ikA_I = -(\beta - ik)B + (\beta + ik)B'$$

$$A_T e^{ika} = \frac{2\beta}{(\beta - ik)} B e^{-\beta a} \quad \text{and} \quad B' = B e^{-2\beta a} \frac{(\beta + ik)}{(\beta - ik)}$$

In the limit of a wide barrier where  $e^{-2\beta a} \ll 1$  we can approximate that  $B \ll B'$ , i.e.  $2ikA_I \approx -(\beta - ik)B$  which gives

$$A_T e^{ika} \approx -\frac{4ik\beta e^{-\beta a}}{(\beta - ik)^2} A_I$$

and

$$T \approx \left[ \frac{16k^2\beta^2}{(\beta^2 + k^2)^2} \right] e^{-2\beta a}$$

Using the definitions

$$k = \frac{\sqrt{2mE}}{\hbar} \quad \text{and} \quad \beta = \frac{\sqrt{2m(V_B - E)}}{\hbar},$$

this can be written as

$$T \approx \left[ \frac{16E(V_B - E)}{V_B^2} \right] e^{-2\beta a}$$

- This is also known as the (QM) *tunneling probability*