

## 1. Proof of Exercise 9 Demo

Show that the solution for the problem:

$$\max_a \left\{ \frac{a^\top B a}{a^\top W a} \right\},$$

is obtained by setting  $a$  equal to the eigenvector of  $W^{-1}B$  that corresponds to the largest eigenvalue.

$W$  = measure of group dispersions       $B$  = dispersion between groups

Since the vector  $a$  can be scaled arbitrarily without affecting the ratio, we can formulate the problem as follows:

$$\max_a \{a^\top B a\} \quad \text{s.t.} \quad a^\top W a = 1.$$

Let  $W^{1/2}$  be the symmetric square root of  $W$ . Note that  $W$  is symmetric. Let  $z = W^{1/2}a$  and  $a = W^{-1/2}z$ . Then

$$\begin{aligned} a^\top B a &= (W^{-1/2}z)^\top B (W^{-1/2}z) = z^\top W^{-1/2} B W^{-1/2} z, \\ a^\top W a &= (W^{-1/2}z)^\top W (W^{-1/2}z) = z^\top z. \end{aligned}$$

Note that  $W^{-1/2} B W^{-1/2}$  is symmetric and hereby the spectral decomposition exists ( $W^{-1/2} B W^{-1/2} = \Gamma \Lambda \Gamma^\top$  and  $\Gamma^\top \Gamma = I$ ),

$$\begin{aligned} z^\top W^{-1/2} B W^{-1/2} z &= z^\top \Gamma \Lambda \Gamma^\top z = w^\top \Lambda w, & (w = \Gamma^\top z) \\ z^\top z &= z^\top \Gamma \Gamma^\top z = w^\top w. \end{aligned}$$

Now we can reformulate the problem as

$$\max_w \{w^\top \Lambda w\} = \max_w \left\{ \sum_{i=1}^p \lambda_i w_i^2 \right\} \quad \text{s.t.} \quad w^\top w = 1.$$

Since  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_p$ , we choose the first element of  $w$  to be one and the rest to be 0. This means that  $z = \Gamma w = \gamma_1$ , where  $\gamma_1$  is the first eigenvector of  $W^{-1/2} B W^{-1/2}$  and  $a = W^{-1/2}z = W^{-1/2}\gamma_1$ .

Note that for any two matrices  $A \in \mathbb{R}^{n \times p}$  and  $C \in \mathbb{R}^{p \times n}$ , the non-zero eigenvalues of  $AC$  and  $CA$  are the same and have the same multiplicity (Theorem A.6.2 of Mardia, Kent and Bibby). Now let  $A = W^{-1/2}B$  and  $C = W^{-1/2}$ , this means that the non-zero eigenvalues of  $CA = W^{-1}B$  are the same as  $AC = W^{-1/2}BW^{-1/2}$ . Hence,  $\lambda_1$  is the largest eigenvalue of  $W^{-1}B$ . Since  $\gamma_1$  is the eigenvector corresponding to the largest eigenvalue  $\lambda_1$  of  $W^{-1/2}BW^{-1/2}$ , we have that

$$\begin{aligned} W^{-1}B(W^{-1/2}\gamma_1) &= W^{-1/2}(W^{-1/2}BW^{-1/2}\gamma_1) = W^{-1/2}\lambda_1\gamma_1 \\ &= \lambda_1(W^{-1/2}\gamma_1). \end{aligned}$$

This shows that  $a = W^{-1/2}\gamma_1$  is the eigenvector of  $W^{-1}B$  corresponding to its largest eigenvalue  $\lambda_1$ .