

Time Series Analysis with R

A. Ian McLeod, Hao Yu, Esam Mahdi

*Department of Statistical and Actuarial Sciences, The University of Western Ontario,
London, Ont., Canada N6A 5B7*

The purpose of our article is to provide a summary of a selection of some of the high-quality published computational time series research using R. A more complete overview of time series software available in R for time series analysis is available in the CRAN¹ task views.² If you are not already an R user, this article may help you in learning about the R phenomenon and motivate you to learn how to use R. Existing R users may find this selective overview of time series software in R of interest. Books and tutorials for learning R are discussed later in this section. An excellent online introduction from the R Development Core Team is available³ as well as extensive contributed documentation.⁴

In the area of computational time series analysis, especially for advanced algorithms, R has established itself as the choice of many researchers. R is widely used not only by researchers but also in diverse time series applications and in the teaching of time series courses at all levels. Naturally, there are many other software systems such as *Mathematica* (Wolfram Research, 2011), that have interesting and useful additional capabilities, such as symbolic computation (Smith and Field, 2001; Zhang and McLeod, 2006). For most researchers working with time series, R provides an excellent broad platform.

The history of R has been discussed elsewhere (Gentleman and Ihaka,

Email addresses: aim@stats.uwo.ca (A. Ian McLeod), hyu@stats.uwo.ca (Hao Yu), emahdi@uwo.ca (Esam Mahdi)

¹Comprehensive R Archive

²<http://cran.r-project.org/web/views/>

³<http://cran.r-project.org/manuals.html>

⁴<http://cran.r-project.org/other-docs.html>

1996) so before continuing our survey we will just point out some other key features of this quantitative programming environment (QPE).

R is an open source project, providing a freely available and a high quality computing environment with thousands of add-on packages. R incorporates many years of previous research in statistical and numerical computing and so it is built on a solid foundation of core statistical and numerical algorithms. The R programming language is a functional, high-level interactive and scripting language that offers two levels of object-oriented programming. For an experienced R user, using this language to express an algorithm is often easier than using ordinary mathematical notation and it is more powerful since, unlike mathematical notation, it can be evaluated. In this way, R is an important tool of thought. Novice and casual users of R may interact with it using Microsoft Excel (Heiberger and Neuwirth, 2009) or R Commander (Fox, 2005).

Through the use of Sweave (Leisch, 2002, 2003), R supports high-quality technical typesetting and reproducible research including reproducible applied statistical and econometric analysis (Kleiber and Zeileis, 2008). This article has been prepared using Sweave and R scripts for all computations, including all figures and tables, are available in an online supplement.⁵ This supplement also includes a PDF preprint of this article showing all graphs in color.

R supports 64-bit, multicore, parallel and cluster computing (Schmidberger et al., 2009; Hoffmann, 2011; Revolution Computing, 2011). Since R is easily interfaced to other programming languages such as C and Fortran, computationally efficient programs may simply be executed in cluster and grid computing environments using R to manage the rather complex message-passing interface.

There is a vast literature available on R that includes introductory books as well as treatments of specialized topics. General purpose introductions to R are available in many books (Braun and Murdoch, 2008; Crawley, 2007; Dalgaard, 2008; Adler, 2009; Everitt and Hothorn, 2009; Zuur et al., 2009). Advanced aspects of the R programming are treated by (Venables and Ripley, 2000; Spector, 2008; Chambers, 2008; Gentleman, 2009). Springer has published more than 30 titles in the *Use R* book series, Chapman &

⁵<http://www.stats.uwo.ca/faculty/aim/tsar.html>

Hall/CRC has many forthcoming titles in *The R Series* and there are many other high quality books that feature R. Many of these books discuss R packages developed by the author of the book and others provide a survey of R tools useful in some application area. In addition to this flood of high quality books, the *Journal of Statistical Software* (JSS) publishes refereed papers discussing statistical software. JSS reviews not only the paper but the quality of the computer code as well and publishes both the paper and code on its website. Many of these papers discuss R packages. The rigorous review process ensures a high quality standard. In this article, our focus will be on R packages that are accompanied by published books and/or papers in JSS.

The specialized refereed journal, *The R Journal*, features articles of interest to the general R community. There is also an interesting BLOG sponsored by Revolution Analytics.⁶

The non-profit association Rmetrics (Würtz, 2004) provides R packages for teaching and research in quantitative finance and time series analysis that are further described in the electronic books that they publish.

There are numerous textbooks, suitable for a variety of courses in time series analysis (Venables and Ripley, 2002; Chan, 2010; Cryer and Chan, 2008; Lütkepohl and Krätzig, 2004; Shumway and Stoffer, 2011; Tsay, 2010). These textbooks incorporate R usage in the book and an R package on CRAN that includes scripts and datasets used in the book.

1. Time series plots

In this section our focus is on plots of time series. Such plots are often the first step in an exploratory analysis and are usually provided in a final report. R can produce a variety of these plots not only for regular time series but also for more specialized time series such as irregularly-spaced time series. The built-in function, `plot()`, may be used to plot simple series such as the annual lynx series, `lynx`. The aspect-ratio is often helpful in visualizing slope changes in a time series (Cleveland et al., 1988; Cleveland, 1993). For many time series an aspect-ratio of 1/4 is good choice. The function `xypplot()` (Sarkar, 2008) allows one to easily control

⁶<http://blog.revolutionanalytics.com/>

the aspect-ratio. Figure 1 shows the time series plot of the lynx series with an aspect-ratio of 1/4. The asymmetric rise and fall of the lynx population is easily noticed with this choice of the aspect-ratio.




Figure 1: Annual numbers of lynx trappings in Canada.

There are many possible styles for your time series plots. Sometimes a high-density line plot is effective as in Figure 2.




Figure 2: High density line plot.

Another capability of `xyplot()` is the cut-and-stack time series plot for longer series. Figure 3 shows a cut-and-stack plot of the famous Beveridge wheat price index using `xyplot()` and `asTheEconomist()`. The cut-and-stack plot uses the equal-count-algorithm (Cleveland, 1993) to divide the series into a specified number of subseries using an overlap. The default setting is for a 50% overlap.

Figure 4 uses `xyplot()` to plot the seasonal decomposition of the well-known CO_2 time series. The seasonal adjustment algorithm available in R `stl()` is described in the R function documentation and in more detail by Cleveland (1993). This plot efficiently reveals a large amount of information. For example, Figure 4, reveals that the seasonal amplitudes are increasing.

index




Figure 3: Beveridge wheat price index.




Figure 4: Atmospheric concentration of CO₂.

Bivariate or multivariate time series may also be plotted with `xyplot()`. In Figure 5, the time series plot for the annual temperature in °C for Canada (CN), Great Britain (UK) and China (CA) 1973-2007, is shown.⁷ Figure 5 uses juxtaposition – each series is in a separate panel. This is often preferable to superposition or showing all series in the same panel. Both types of positioning are available using the R functions `plot()` or `xyplot()`.




Figure 5: Average annual temperature, °C, 1973-2007 for Canada (CN), Great Britain (UK) and China (CN).

A specialized plot for bivariate time series called the cave plot (Becker et al., 1994) is easily constructed in R as shown by Zhou and Braun (2010). When there are many multivariate time series, using `xyplot` may not be feasible. In this case, `mvtsplot()` provided by Peng (2008) may be used. Many interesting examples, including a stock market portfolio, daily time series of ozone pollution in 100 US counties, and levels of sulfate in 98 US counties are discussed by Peng (2008).

⁷The data were obtained from *Mathematica*'s curated databases.

Usually this plot is used with many time series – at least ten or more – but for simplicity and in order to compare with the last example, Figure 6 displays the annual temperature series for Canada, Great Britain and China using `mvtsplot()`. The right panel of the plot shows a boxplot for the values in each series. From this panel it is clear that China is generally much warmer than Great Britain and Canada and that Great Britain is often slightly cooler than Canada on an average annual basis. The bottom panel shows the average of the three series. The image shown shows the variation in the three series. The colors purple, grey and green correspond to low, medium and high values for each series. The darker the shading the larger the value. From image in Figure 6, it is seen that Canada has experienced relatively warmer years than Great Britain or China since about the year 2000. During 1989 to 1991 the average annual temperature in Canada was relatively low compared to Great Britain and China. There are many more possible option choices for constructing these plots (Peng, 2008). This plot is most useful for displaying a large number of time series.




Figure 6: Average annual temperature in °C, 1973-2007.

Financial time series are often observed on a daily basis but not including holidays and other days when the exchange is closed. Historical and current stock market data may be accessed using `get.hist.quote()` (Trapletti, 2011). Dealing with dates and times is often an important practical issue with financial time series. Grolemond and Wickham (2011) provide a new approach to this problem and review the other approaches that have been used in R. Irregularly observed time series can be plotted using Rmetrics functions (Wuertz and Chalabi, 2011). The RMetrics package **flmport** also has functions for retrieving stock market data from various stock exchanges around the world.

In Figure 7, the function `yahooSeries()` is used to obtain the last 60 trading days of the close price of IBM stock. The function RMetrics `timeSeries()` converts this data to a format that can be plotted.




Figure 7: IBM, daily close price, returns in percent.

Time series plots are ubiquitous and important in time series applications. It must also be noted that R provides excellent time series graphic capabilities with other standard time series functions, including functions time series diagnostics, autocorrelations, spectral analysis, and

wavelet decompositions to name a few. The output from such functions is usually best understood from the graphical output.

More generally, there are many other types of functions available for data visualization and statistical graphics. For example, all figures in the celebrated monograph on visualizing data by [Cleveland \(1993\)](#) may be reproduced using the R scripts.⁸

The R package `ggplot2` ([Wickham, 2009](#)) implements the novel graphical methods discussed in the wonderful graphics book by ([Wilkinson, 1999](#)). An interesting rendition of Millard's famous temporal-spatial graph of Napoleon's invasion of Russia using **ggplot2** is available in the online documentation.

Dynamic data visualization, including time series, is provided with **rggobi** ([Cook and Swayne, 2007](#)).

The foundation and the state-of-the-art in R graphics is presented in the book by [Murrell \(2011\)](#).

2. Base packages: stats and datasets

The **datasets** and **stats** packages are normally automatically loaded by default when R is started. These packages provide a comprehensive suite of functions for analyzing time series, as well as many interesting time series datasets. These datasets are briefly summarized in the Appendix (§12.1).

The **stats** package provides the base functions for time series analysis. These functions are listed in the Appendix (12.2). For further discussion of these functions, see [Cowpertwait and Metcalfe \(2009\)](#). Many time series textbooks provide a brief introduction to R and its use for time series analysis ([Cryer and Chan, 2008](#); [Shumway and Stoffer, 2011](#); [Venables and Ripley, 2002](#); [Wuertz, 2010](#)).

[Adler \(2009\)](#) provides a comprehensive introduction to R that includes a chapter on time series analysis.

An introduction to ARIMA models and spectral analysis with R is given in the graduate level applied statistics textbook by [Venables and Ripley \(2002\)](#). This textbook is accompanied by the R package **MASS**.

⁸<http://www.stat.purdue.edu/~wsc/visualizing.html>

The time series analysis functions that R provides are sufficient to supplement most textbooks on time series analysis.

2.1. **stats**

First we discuss the **stats** time series functions. In addition to many functions for manipulating time series such as filtering, differencing, inverse differencing, windowing, simulating, aggregating and forming multivariate series, there is a complete set of functions for auto/cross correlations analysis, seasonal decomposition using moving-average filters or loess, univariate and multivariate spectral analysis, univariate and multivariate autoregression, and univariate ARIMA model fitting. Many of these functions implement state-of-the art algorithms. The `ar()` function includes options, in both the univariate and multivariate cases, for Yule-Walker, least-squares or Burg estimates. Although `ar()` implements the maximum likelihood estimator, the package **FitAR** (McLeod et al., 2011b; McLeod and Zhang, 2008b) provides a faster and more reliable algorithm.

The function `spectrum()`, also for both univariate and multivariate series, implements the iterated Daniel smoother (Bloomfield, 2000) and in the univariate case, the autoregressive spectral density estimator (Percival and Walden, 1993).

The `arma()` function implements a Kalman filter algorithm that provides exact maximum likelihood estimation and an exact treatment for the missing-values (Ripley, 2002). This function is interfaced to C code to provide maximum computational efficiency. The `arma()` function has options for multiplicative seasonal ARIMA model fitting, subset models where some parameters are fixed at zero, and regression with ARIMA errors. The functions `tsdiag()` and `Box.test()` provide model diagnostic checks. For ARMA models, a new maximum likelihood algorithm (McLeod and Zhang, 2008a) written entirely in the R language is available in the **FitARMA** package (McLeod, 2010).

A brief example of a medical intervention analysis carried out using `arma()` will now be discussed. In a medical time series of monthly average creatinine clearances, a step intervention analysis model with a multiplicative seasonal ARIMA(0, 1, 1) (1, 0, 0)₁₂ error term was fit. The intervention effect was found to be significant at 1%. To illustrate this finding, Figure 8 compares the forecasts before and after the intervention

date. The forecasts are from a model fit to the pre-intervention series. The plot visually confirms the decrease in creatinine clearances after the intervention.




Figure 8: Creatinine clearance series.

Exponential smoothing methods are widely used for forecasting (Gelper et al., 2010) and are available in `stats` (Meyer, 2002). Simple exponential smoothing defines the prediction for z_{t+h} , $h = 1, 2, \dots$ as \hat{z}_{t+1} where $\hat{z}_{t+1} = \lambda z_t + (1 - \lambda)\hat{z}_{t-1}$. The forecast with this method is equivalent to that from an ARIMA(0,1,1). An extension, double exponential smoothing, forecasts z_{t+h} , $h = 1, 2, \dots$ uses the equation $\hat{z}_{t+h} = \hat{a}_t + h\hat{b}_t$, where $\hat{a}_t = \alpha z_t + (1 - \alpha)(\hat{a}_{t-1} + \hat{b}_{t-1})$, $\hat{b}_t = \beta(\hat{a}_t - \hat{a}_{t-1}) + (1 - \beta)\hat{b}_{t-1}$, where α and β are the smoothing parameters. Double exponential smoothing is sometimes called Holt's linear trend method and it can be shown to produce forecasts equivalent to the ARIMA(0,2,2). The Winter's method for seasonal time series with period p , forecasts z_{t+h} , by $\hat{z}_{t+h} = \hat{a}_t + h\hat{b}_t + \hat{s}_t$, where $\hat{a}_t = \alpha(z_t - \hat{s}_{t-p}) + (1 - \alpha)(\hat{a}_{t-1} + \hat{b}_{t-1})$, $\hat{b}_t = \beta(\hat{a}_t - \hat{a}_{t-1}) + (1 - \beta)\hat{b}_{t-1}$, $\hat{s}_t = \gamma(Y - \hat{a}_t) + (1 - \gamma)\hat{s}_{t-p}$, α , β and γ are smoothing parameters. In the multiplicative version, $\hat{z}_{t+h} = (\hat{a}_t + h\hat{b}_t)\hat{s}_t$. Winter's method is equivalent to the multiplicative seasonal ARIMA airline-model in the linear case. All of the above exponential smoothing models may be fit with `HoltWinters()`. This function also has `predict()` and `plot()` methods.

Structural time series models (Harvey, 1989) are also implemented using

Kalman filtering in the function `StructTS()`. Since the Kalman filter is used, Kalman smoothing is also available and it is implemented in the function `tsSmooth()`. The basic structural model is comprised of an observational equation,

$$z_t = \mu_t + s_t + e_t, \quad e_t \sim NID(0, \sigma_e^2)$$

and the state equations,

$$\mu_{t+1} = \mu_t + \xi_t, \quad \xi_t \sim NID(0, \sigma_\xi^2),$$

$$v_{t+1} = v_t + \zeta_t, \quad \zeta_t \sim NID(0, \sigma_\zeta^2),$$

$$\gamma_{t+1} = -(\gamma_t + \dots + \gamma_{t-s+2}) + \omega_t, \quad \omega_t \sim NID(0, \sigma_\eta^2).$$

If σ_ω^2 is set to zero, the seasonality is deterministic. The local linear trend model is obtained by omitting the term involving γ_t in the observational equation and the last state equation may be dropped as well. Setting $\sigma_\xi^2 = 0$ in the local linear trend model results in a model equivalent to the ARIMA(0,2,2). Setting $\sigma_\xi^2 = 0$ produces the local linear model which is also equivalent to the ARMA(0,1,1).

In Figure 9, the forecasts from the multiplicative Winter’s method for the next 12 months are compared with forecasts from the multiplicative-seasonal ARIMA(0, 1, 1) (0, 1, 1)₁₂ model. With this model, logarithms of the original data were used and then the forecasts were back-transformed. There are two types of backtransform that may be used for obtaining the forecasts in the original data domain (Granger and Newbold, 1976; Hopwood et al., 1984) — naive or minimum-mean-square-error (MMSE). Figure 9 compares these backtransformed forecasts and shows that the MMSE are shrunk relative to the naive forecasts.

2.2. tseries

The `tseries` package (Trapletti, 2011) is well-established and provides both useful time series functions and datasets. These are summarized in Appendix (12.3).




Figure 9: Comparisons of forecasts for 1961.

2.3. Forecast

The package **Forecast** (Hyndman, 2010) provides further support for forecasting using ARIMA and a wide class of exponential smoothing models. These methods are described briefly by Hyndman and Khandakar (2008) and in more depth in the book (Hyndman et al., 2008). Hyndman and Khandakar (2008) discuss a family of sixty different exponential smoothing models and provide a new state-space approach to evaluate the likelihood function.

Appendix 12.4, Table 16 summarizes functions for exponential smoothing models.

Automatic ARIMA and related functions are summarized in Table 15.

In addition, general utility functions that are useful for dealing with time series data such as number of days in each month, interpolation for missing values, a new seasonal plot, and others are briefly described in Table 14.

3. More Linear Time Series Analysis

3.1. State space models and Kalman filtering

Tusell (2011) provides an overview of Kalman filtering with R. In addition to **StructTS**, there are four other packages that support Kalman

filtering and state-space modeling of time series. In general, the state space model (Harvey, 1989; Tusell, 2011) is comprised of two equations, the observation equation:

$$\mathbf{y}_t = \mathbf{d}_t + \mathbf{Z}_t \boldsymbol{\alpha}_t + \boldsymbol{\epsilon}_t \quad (1)$$

and the state equation:

$$\boldsymbol{\alpha}_t = \mathbf{c}_t + \mathbf{T}_t \boldsymbol{\alpha}_{t-1} + \mathbf{R}_t \boldsymbol{\eta}_t, \quad (2)$$

where the white noises, $\boldsymbol{\epsilon}_t$ and $\boldsymbol{\eta}_t$, are multivariate normal with mean vector zero and covariance matrices \mathbf{Q}_t and \mathbf{H}_t respectively. The white noise terms are uncorrelated, $E\{\boldsymbol{\epsilon}'_t \boldsymbol{\eta}_t\} = 0$.

The Kalman filter algorithm recursively computes,

- predictions for $\boldsymbol{\alpha}_t$
- predictions for \mathbf{y}_t
- interpolation for \mathbf{y}_t

and in each case the estimated covariance matrix is also obtained.

Dropping the terms \mathbf{d}_t and \mathbf{c}_t and restricting all the matrices to be constant over time provides a class of state-space models that includes univariate and multivariate ARMA models (Brockwell and Davis, 1991; Gilbert, 1993; Durbin and Koopman, 2001). As previously mentioned, the built-in function `arma` uses a Kalman filter algorithm to provide exact MLE for univariate ARIMA with missing values (Ripley, 2002). The `dse` package Gilbert (2011) implements Kalman filtering for the time-invariant case and provides a general class of models that includes multivariate ARMA and ARMAX models.

Harrison and West (1997) and Harvey (1989) provide a comprehensive account of Bayesian analysis dynamic linear models based on the Kalman filter and this theme is further developed in the book by Petris et al. (2009). This book also provides illustrative R scripts and code. The accompanying package `dlm` (Petris, 2010) provides functions for estimation and filtering as well as a well-written vignette explaining how to use the software.

The following example of fitting the random walk plus noise model,

$$\begin{aligned} y_t &= \theta_t + v_t, & v_t &\sim \mathcal{N}(0, V) \\ \theta_t &= \theta_{t-1} + w_t, & w_t &\sim \mathcal{N}(0, W) \end{aligned}$$

to the Nile series and plotting the filtered series, Figure 10 and its 95% interval, is taken from the vignette by [Petris \(2010\)](#).




Figure 10: Nile river flows (solid line with circles), filter values after fitting random walk with noise (solid thick line) and 95% confidence interval (dashed lines).

Three other packages for Kalman filtering ([Dethlefsen et al., 2009](#); [Luethi et al., 2010](#); [Helske, 2011](#)) are also reviewed by [Tusell \(2011\)](#).

3.2. An approach to linear time series analysis using Durbin-Levinson recursions

Table 17 in Appendix 12.5 lists the main functions available in the **ltsa** package for linear time series analysis.

The Durbin-Levinson recursions ([Box et al., 2008](#)) provide a simple and direct approach to the computation of the likelihood, computation of exact forecasts and their covariance matrix, and simulation for any linear process

defined by its autocorrelation function. This approach is implemented in **ltsa** (McLeod et al., 2007, 2011a).

In Section 3.3, this approach is implemented for the fractional Gaussian noise (FGN) and a comprehensive model building R package is provided for this purpose using the functions in **ltsa**.

Three methods of simulating a time series given its autocovariance function are available: `DHSimulate()`, `DLSimulate()`, and `SimGLP()`. `DHSimulate()` implements the fast Fourier algorithm (FFT) of Davies and Harte (1987). But this algorithm is not applicable for all stationary series (Craigmile, 2003) so `DHSimulate()`, based on the Durbin-Levinson recursion, is also provided. The algorithm `SimGLP()` is provided for simulating a time series with non-Gaussian innovations based on the equation,

$$z_t = \mu + \sum_{i=1}^{\varrho} \psi_i a_{t-i}. \quad (3)$$

The sum involved in Equation (3) is efficiently evaluated using the R function `convolve()` that uses the fast Fourier transform (FFT) method. The built-in function `arima.sim()` may also be used in the case of ARIMA models. The functions `TrenchInverse()` and `TrenchInverseUpdate()` are useful in some applications involving Toeplitz covariance matrices. `TrenchForecast()` provides exact forecasts and their covariance matrix.

The following illustration is often useful in time series lectures when forecasting is discussed. In the next example we fit an AR(9) to the annual sunspot numbers, 1700-1988, `sunspot.year`. For forecasting computations, it is standard practice to treat the parameters as known, that is to ignore the error due to estimation. This is reasonable because the estimation error is small in comparison to the innovations. This assumption is made in our algorithm `TrenchForecast()`. Letting $z_m(\ell)$ denote the optimal minimum mean square error forecast at origin time $t = m$ and lead time ℓ , we compare the forecasts of z_{m+1}, \dots, z_n using the one-step ahead predictor $z_{m+\ell-1}(1)$, with the fixed origin prediction $z_m(\ell)$, where $\ell = 1, \dots, L$ and $L = n - m + 1$. Figure 11 compares forecasts and we see many interesting features. The fixed origin forecasts are less accurate as might be expected. As well the fixed origin forecasts show systematic departures whereas the one-step do not.




Figure 11: Comparing forecasts from a fixed origin, 1969, with lead-one forecasts starting in 1969 for `sunspot.year`.

As shown by this example, `TrenchForecast()` provides a more flexible approach to forecasting than provided by `predict()`.

3.3. Long memory time series analysis

Let z_t , $t = 1, 2, \dots$ be stationary with mean zero and autocovariance function, $\gamma_z(k) = \text{cov}(z_t, z_{t-k})$. Many long memory processes such as the FGN (fractional Gaussian Noise) and FARMA (fractional ARMA) may be characterized by the property that $k^\alpha \gamma_z(k) \rightarrow c_{\alpha, \gamma}$ as $k \rightarrow \infty$, for some $\alpha \in (0, 1)$ and $c_{\alpha, \gamma} > 0$. Equivalently,

$$\gamma_z(k) \sim c_{\alpha, \gamma} k^{-\alpha}.$$

The FARMA and FGN models are reviewed by [Hipel and McLeod \(1994\)](#); [Beran \(1994\)](#); [Brockwell and Davis \(1991\)](#). FGN can simply be described as a stationary Gaussian time series with covariance function, $\rho_k = (|k+1|^{2H} - 2|k|^{2H} + |k-1|^{2H})/2$, $0 < H < 1$. The FARMA model generalizes the ARIMA model to a family of stationary models with fractional difference parameter d , $d \in (-0.5, 0.5)$. The long-memory parameters H and d may be expressed in terms of α ,

$H \simeq 1 - \alpha/2$, $H \in (0, 1)$, $H \neq 1/2$ and $d \simeq 1/2 - \alpha/2$, $d \in (-1/2, 1/2)$, $d \neq 0$ (McLeod, 1998). Gaussian white noise corresponds to $H = 1/2$ and in the case of FARMA, $d = 0$ assuming no AR or MA components. Haslett and Raftery (1989) developed an algorithm for maximum likelihood estimation of FARMA models and applied these models to the analysis of long wind speed time series. This algorithm is available in R in the package `fracdiff` (Fraley et al., 2009). The generalization of the FARMA model to allow more general values of d is usually denoted by ARFIMA. A frequently cited example of a long-memory time series is the minimum annual flows of the Nile over the period 622-1284, $n = 663$ (Percival and Walden, 2000, §9.8). The package `longmemo` (Beran et al., 2009) has this data as well as other time series examples. **FGN** provides exact MLE for the parameter H as well as a parametric bootstrap and minimum mean square error forecast. For the Nile data, $\hat{H} = 0.831$. The time series plots in Figure 12 show the actual Nile series along with three bootstraps.




Figure 12: Comparing actual Nile minima series with two bootstrap versions.

As a further illustration of the capabilities of R, a simulation experiment was done to compare the estimation of the H -parameter in fractional Gaussian noise using the exact MLE function `FitFGN()` in **FGN** and the GLM method `FEXPEst()` in the package `longmemo`. The function `SimulateFGN()` in **FGN** was used to simulate 100 sequences of length $n = 200$ for $H = 0.3, 0.5, 0.7$. Each sequence was fit by the MLE and GLM method and the absolute error of the difference between the estimate and

the true parameter was obtained, that is, $\text{Err}_{\text{MLE}} = |\hat{\mathbf{H}}_{\text{MLE}} - \mathbf{H}|$ and $\text{Err}_{\text{GLM}} = |\hat{\mathbf{H}}_{\text{GLM}} - \mathbf{H}|$. From Figure 13, the notched boxplot for $\text{Err}_{(\text{GLM})} - \text{Err}_{(\text{MLE})}$, we see that the MLE is more accurate. These computations take less than 30 seconds using direct sequential evaluation on a current PC.




Figure 13: Comparing MLE estimator and GLM estimator for the parameter H in fractional Gaussian noise.

The ARFIMA model extends the FARMA models to the ARIMA or difference-stationary case (Diebold and Rudebusch, 1989; Baillie, 1996). The simplest approach is to choose the differencing parameter and then fit the FARMA model to the differenced time series.

3.4. Subset autoregression

The **FitAR** package (McLeod and Zhang, 2006, 2008b; McLeod et al., 2011b) provides a more efficient and reliable exact MLE for AR(p) than is available with the built-in function `ar()`. Two types of subset autoregressions may also be fit. The usual subset autoregression may be written, $\phi(\mathbf{B})(z_t - \mu) = a_t$, where $\phi(\mathbf{B}) = 1 - \phi_{i_1}B - \dots - \phi_{i_m}B^{i_m}$, where i_1, \dots, i_m are the subset of lags. For this model, ordinary least squares (OLS) is used to estimate the parameters. The other subset family is parameterized using

the partial autocorrelations as parameters. Efficient model selection, estimation and diagnostic checking algorithms are discussed by [McLeod and Zhang \(2006\)](#) and [McLeod and Zhang \(2008b\)](#) and implemented in the **FitAR** package ([McLeod et al., 2011b](#)). Any stationary time series can be approximated by a high order autoregression that may be selected using one of several information criteria. Using this approximation, **FitAR**, provides functions for automatic bootstrapping, spectral density estimation, and Box-Cox analysis for any time series. The optimal Box-Cox transformation for the `lynx` is obtained simply from the command `R > BoxCox(lynx)`. The resulting plot is shown in [Figure 14](#).




Figure 14: Box-Cox analysis of lynx time series.

The functions of interest in the **FitAR** package are listed in [Appendix 12.6](#).

3.5. Periodic autoregression

Let z_t , $t = 1, \dots, n$ be n consecutive observations of a seasonal time series with seasonal period s . For simplicity of notation, assume that $n/s = N$ is an integer, so N full years of data are available. The time index parameter, t , may be written $t = t(r, m) = (r - 1)s + m$, where $r = 1, \dots, N$

and $m = 1, \dots, s$. In the case of monthly data, $s = 12$ and r and m denote the year and month. If the expected monthly mean $\mu_m = E\{z_{t(r,m)}\}$ and the covariance function, $\gamma_{\ell,m} = \text{cov}(z_{t(r,m)}, z_{t(r,m)-\ell})$ depend only on ℓ and m , z_t is said to be periodically autocorrelated and is periodic stationary. The periodic AR model of order (p_1, \dots, p_s) may be written,

$$z_{t(r,m)} = \mu_m + \sum_{i=1}^{p_m} \phi_{i,m}(z_{t(r,m)-i} - \mu_{m-i}) + a_{t(r,m)}, \quad (1.3)$$

where $a_{t(r,m)} \sim \text{NID}(0, \sigma_m^2)$, where m obeys modular arithmetic base s . This model originated in monthly streamflow simulation and is further discussed with examples by [Hipel and McLeod \(1994\)](#). Diagnostic checks for periodic autoregression are derived by [McLeod \(1994\)](#). The package `pear` ([McLeod and Balcilar, 2011](#)) implements functions for model identification, estimation and diagnostic checking for periodic AR models.

We conclude with a brief mention of some recent work on periodically correlated time series models which we hope to see implemented in R. [Tsfaye et al. \(2011\)](#) develop a parsimonious and efficient procedure for dealing with periodically correlated daily ARMA series and provide applications to geophysical series. [Ursu and Duchesne \(2009\)](#) extend modeling procedures to the vector PAR model and provide an application to macro economic series. [Aknouche and Bibi \(2009\)](#) show that quasi-MLE provide consistent, asymptotically normal estimates in a periodic GARCH model under mild regularity conditions.

4. Time series regression

An overview of selected time series regression topics is given in this section. Further discussion of these and other topics involving time series regression with R is available in several textbooks ([Cowpertwait and Metcalfe, 2009](#); [Cryer and Chan, 2008](#); [Kleiber and Zeileis, 2008](#); [Shumway and Stoffer, 2011](#)).

4.1. Cigarette consumption data

Most of the regression methods discussed in this section will be illustrated with data from an empirical demand analysis for cigarettes in

Canada ([Thompson and McLeod, 1976](#)). The variables of interest, consumption of cigarettes per capita, Q_t , real disposable income per capita, Y_t , and the real price of cigarettes, P_t , for $t = 1, \dots, 23$ corresponding to the years 1953-1975 were all logarithmically transformed and converted to an R dataframe `cig`. For some modeling purposes, it is more convenient to use a `ts` object,

```
R >cig.ts <- ts(as.matrix.data.frame(cig), start = 1953,
+             freq = 1)
```

The time series are shown in Figure 15.

```
R >plot(cig.ts, xlab = "year", main = "", type = "o")
```




Figure 15: Canadian cigarette data, consumption/adult(Q), real price(P), income/adult(Y).

4.2. Durbin-Watson test

The exact p-value for the Durbin-Watson diagnostic test for lack of autocorrelation in a linear regression with exogenous inputs and Gaussian white noise errors is available with the function `dwtest()` in the **lmtest** package ([Hothorn et al., 2010](#)). The diagnostic check statistic may be written

$$d = \frac{\sum_{t=2}^n (\hat{e}_t - \hat{e}_{t-1})^2}{\sum_{t=1}^n \hat{e}_t^2}, \quad (4)$$

where $\hat{e}_t, t = 1, \dots, n$ are the OLS residuals. Under the null hypothesis, d should be close to 2 and small values of d indicate positive autocorrelation.

Many econometric textbooks provide tables for the critical values of d . But in small samples these tables may be inadequate since there is a fairly large interval of values for d for which the test is inconclusive. This does not happen when the exact p-value is computed. Additionally, current statistical practice favors reporting p-values in diagnostic checks (Moore, 2007).

The Durbin-Watson test is very useful in time series regression for model selection. When residual autocorrelation is detected, sometimes simply taking first or second differences is all that is needed to remove the effect of autocorrelation. In the next example we find that taking second differences provides an adequate model.

First we fit the empirical demand equation, regressing demand Q_t on real price P_t and income Y_t , $Q_t = \beta_0 + \beta_1 P_t + \beta_2 Y_t + e_t$ using OLS with the `lm()` function. Some of the output is shown below.

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	3.328610	2.5745756	1.2928771	2.107900e-01
P	-0.402811	0.4762785	-0.8457468	4.076991e-01
Y	0.802143	0.1118094	7.1741970	6.011946e-07

This output suggests P_t is not significant but Y_t appears to be highly significant. However, since the Durbin-Watson test rejects the null hypothesis of no autocorrelation, these statistical inferences about the coefficients in the regression are incorrect.

After differencing, the Durbin-Watson test still detects significant positive autocorrelation.

Finally, fitting the model with second-order differencing, $\nabla^2 Q_t = \beta_0 + \nabla^2 \beta_1 P_t + \nabla^2 \beta_2 Q_t + e_t$, $\hat{\beta}_1 = 0.557$ with a 95% margin of error, 0.464, so the price elasticity is significant at 5%. As may be seen for the computations reproduced below the other parameters are not statistically significant at 5%.

```
R >cig2.lm <- lm(Q ~ P + Y, data = diff(cig.ts, differences = 2))
R >summary(cig2.lm)$coefficients
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-0.003118939	0.008232764	-0.3788447	0.70923480
P	-0.557623890	0.236867207	-2.3541625	0.03012373
Y	0.094773991	0.278979070	0.3397172	0.73800132

The intercept term, corresponds to a quadratic trend, is not significant and can be dropped. Income, Y_t is also not significant. The evidence for lag-one autocorrelation is not strong,

```
R >dwtest(cig2.lm, alternative = "two.sided")
```

Durbin-Watson test

```
data: cig2.lm
DW = 2.6941, p-value = 0.08025
alternative hypothesis: true autocorelation is not 0
```

There is also no evidence of non-normality using the Jarque-Bera test. We use the function `jarque.bera.test()` in the `tseries` package ([Trapletti, 2011](#)).

```
R >jarque.bera.test(resid(cig2.lm))
```

Jarque Bera Test

```
data: resid(cig2.lm)
X-squared = 1.1992, df = 2, p-value = 0.549
```

[Kleiber and Zeileis \(2008, §7\)](#) discuss lagged regression models for time series. and present illustrative simulation experiment using R that compares the power of the Durbin-Watson test with the Breusch-Godfrey test for detecting residual autocorrelation in time series regression ([Kleiber and Zeileis, 2008, §7.1](#)).

As discussed below in Section 4.4, fitting regression with lagged inputs is best done using the package **dynlm**.

4.3. Regression with autocorrelated error

The built-in function `arima` can fit the linear regression model with k inputs and $\text{ARIMA}(p, d, q)$ errors, $y_t = \beta_0 + \beta_1 x_{1,t} + \dots + \beta_k x_{k,t} + e_t$, where $e_t \sim \text{ARIMA}(p, d, q)$ and $t = 1, \dots, n$.

We illustrate by fitting an alternative to the regression just fit above for the Canadian cigarette data.

```
R >with(cig, arima(Q, order = c(1, 1, 1), xreg = cbind(P,
+      Y)))
```

Call:

```
arima(x = Q, order = c(1, 1, 1), xreg = cbind(P, Y))
```

Coefficients:

	ar1	ma1	P	Y
	0.9332	-0.6084	-0.6718	0.2988
s.e.	0.1010	0.2007	0.2037	0.2377

sigma^2 estimated as 0.0008075: log likelihood = 46.71, aic = -83.41

This model agrees well with the linear regression using second differencing.

4.4. Regression with lagged variables

Linear regression models with lagged dependent and/or independent variables are easily fit using the `dynlm` package (Zeileis, 2010). In the case of the empirical demand for cigarettes, it is natural to consider the possible effect lagged price. $\nabla^2 Q_t = \beta_1 \nabla^2 P_t + \beta_{1,2} \nabla^2 P_{t-1} + \beta_2 \nabla^2 Y_t + e_t$,

```
R >summary(dynlm(Q ~ -1 + P + L(P) + Y, data = diff(cig.ts,
+      differences = 2)))$coefficients
```

	Estimate	Std. Error	t value	Pr(> t)
P	-0.6421079	0.2308323	-2.7817077	0.01278799
L(P)	-0.1992065	0.2418089	-0.8238177	0.42145104
Y	-0.2102738	0.2993858	-0.7023507	0.49196623

We see that lagged price is not significant.

4.5. Structural Change

[Brown et al. \(1975\)](#) introduced recursive residuals and related methods for examining graphically the stability of regression over time. These methods and recent developments in testing and visualizing structural change in time series regression are discussed in the book by [Kleiber and Zeileis \(2008, §6.4\)](#) and implemented in the package **strucchange** ([Zeileis et al., 2010, 2002](#)). We use a CUMSUM plot of the recursive residuals to check the regression using second differences for stability. No instability is detected with this analysis.




Figure 16: Cusum test of residuals in cigarette demand regression.

4.6. Generalized linear models

[Kedem and Fokianos \(2002\)](#) provide a mathematical treatment of the use of generalized linear models (GLM) for modeling stationary binary, categorical and count time series. GLM models can account for autocorrelation by using lagged values of the dependent variable in the systematic component. Under regularity conditions, inferences based on large sample theory for GLM time series models can be made using standard software for fitting regular GLM models ([Kedem and Fokianos, 2002, §1.4](#)). In R, the function `glm()` may be used and it is easy to verify

estimates of the precision using the `boot()` function. These GLM-based time series models are extensively used with longitudinal time series (Li, 1994).

As an illustration, we consider the late night fatality data discussed in Vingilis et al. (2005). The purpose of this analysis was to investigate the effect of the extension of bar closing hours to 2:00 AM that was implemented May 1, 1996. This type of intervention analysis (Box and Tiao, 1975) is known as an interrupted time series design in the social sciences (Shadish et al., 2001). The total fatalities per month for the period starting January 1992 and through to December 1999, corresponding to a time series of length $n = 84$, are shown in Figure 17.




Figure 17: Late night car fatalities in Ontario. Bar closing hours were extended May 1996.

The output from the `glm()` function using `y` as the dependent variable, `y1` as the lagged dependent variable⁹, and `x` as the step intervention defined as 0 before May 1, 1996 and 1 after.

```
R >summary(ans)$coefficients
```

⁹ `y` and `y1` are the vectors containing the sequence of observed fatalities and its lagged values.

	Estimate	Std. Error	z value	Pr(> z)
(Intercept)	-2.53923499	0.5040873	-5.03729193	4.721644e-07
x2	1.16691417	0.6172375	1.89054329	5.868534e-02
y1	-0.06616152	0.6937560	-0.09536712	9.240232e-01

The resulting GLM model may be summarized as follows. The total fatalities per month, y_t , are Poisson distributed with mean μ_t , where $\hat{\mu}_t = \exp\{\hat{\beta}_0 + \hat{\beta}_1 x_t + \hat{\beta}_2 y_{t-1}\}$, $\hat{\beta}_0 \doteq -2.54$, $\hat{\beta}_1 \doteq 1.17$, and $\hat{\beta}_2 \doteq -0.07$. There is no evidence of lagged dependence but the intervention effect, β_2 is significant with $p < 0.10$.

We verified the standard deviation estimates of the parameters by using a non-parametric bootstrap with 1000 bootstrap samples. This computation takes less than 10 seconds on most current PC's. Table 1, produced directly from the R output using the package **xtable**, compares the asymptotic and bootstrap standard deviations. As seen from the table the agreement between the two methods is reasonably good.

	(Intercept)	x2	y1
asymptotic	0.50	0.62	0.69
bootstrap	0.49	0.66	0.75

Table 1: Comparison of asymptotic and bootstrap estimates of the standard deviations in the GLM time series regression

Hidden Markov models provide another time series generalization of Poisson and binomial GLM models ([Zucchini and MacDonald, 2009](#)).

5. Nonlinear time series models

Volatility models including the GARCH family of models are one of the newest types on nonlinear time series models. Nonlinear regression models can sometimes be applied to time series. GLM models provide an extension of linear models that is useful for modeling logistic and count time series ([Kedem and Fokianos, 2002](#)). [Ritz and Streibig \(2008\)](#) provides an overview of nonlinear regression models using R. Loess regression in R provides a flexible nonparametric regression approach to handling up to three inputs. Using generalized additive models (GAM), many more inputs could be

accommodated (Wood, 2006). Two packages, **earth** (Milborrow, 2011) and **mda** (Hastie and Tibshirani, 2011) implement MARS or multiadaptive regression splines (Friedman, 1991). Lewis and Stevens (1991) reported that MARS regression produced better out-of-sample forecasts for the the annual sunspot series than competing nonlinear models. In the remainder of the section we discuss tests for nonlinearity and two popular approaches to modeling and forecasting nonlinear time series, threshold autoregression, and neural net.

5.1. Tests for nonlinear time series

One approach is to fit a suitable ARIMA or other linear time series model and then apply the usual Ljung-Box portmanteau test to the squares of the residuals. McLeod and Li (1983) suggested this as a general test for nonlinearity. The built-in function `Box.test()` provides a convenient function for performing this test. Two tests (Terasvirta et al., 1993; Lee et al., 1993) for neglected nonlinearity that are based on neural nets are implemented in **tseries** (Trapletti, 2011) as functions `terasvirta.test()` and `white.test()`. The Keenan test for nonlinearity (Keenan, 1985) is available in **TSA** (Chan, 2011) and is discussed in the textbook by Cryer and Chan (2008).

5.2. Threshold models

Threshold autoregression (TAR) provides a general flexible family for nonlinear time series modeling that has proved useful in many applications. This approach is well suited to time series with stochastic cyclic effects such as exhibited in the annual sunspots or lynx time series. The model equation for a two-regime TAR model may be written,

$$y_t = \phi_{1,0} + \phi_{1,1}y_{t-1} + \dots + \phi_{1,p}y_{t-p} + I(y_{t-d} > r)\{\phi_{2,0} + \phi_{2,1}y_{t-1} + \dots + \phi_{2,p}y_{t-p}\} + \sigma a_t \quad (5)$$

where $I(y_{t-d} > r)$ indicates if $y_{t-d} > r$ the result is 1 and otherwise it is 0. The parameter d is the delay parameter and r is the threshold. There are separate autoregression parameters for each regime. This model may be estimated by least squares or more generally using conditional maximum likelihood.

A TAR model for the predator time series in Figure 18 is described in the book by Cryer and Chan (2008). The package **TSA** (Chan, 2011) provides illustrative datasets from the book (Cryer and Chan, 2008) as well as the function `tar()` for fitting two regime TAR models, methods functions `predict()` and `tsdiag()`, and functions `tar.skelton()` and `tar.sim()`.




Figure 18: Number of prey individuals (*Didinium natsutum*) per ml measured every twelve hours over a period of 35 days.

TAR and related models are also discussed by Tsay (2010) and some R scripts are provided as well the companion package **FinTS** (Graves, 2011) that includes data sets from the book. Figure 19 shows monthly U.S. unemployment. Tsay (2010, Example 4.2) fits the two regime TAR model,

$$\begin{aligned}
 y_t &= 0.083y_{t-2} + 0.158y_{t-3} + 0.0118y_{t-4} - 0.180y_{t-12} + a_{1,t} & \text{if } y_{t-1} \leq 0.01, \\
 &= 0.421y_{t-2} + 0.239y_{t-3} - 0.127y_{t-12} + a_{2,t} & \text{if } y_{t-1} > 0.01,
 \end{aligned}$$

where y_t is the differenced unemployment series. The estimated standard deviations of $a_{1,t}$ and $a_{2,t}$ were 0.180 and 0.217. Tsay (2010) remarks that the TAR provides more insight into the time-varying dynamics of the unemployment rate than the ARIMA.




Figure 19: U.S. civilian unemployment rate, seasonally adjusted, January 1948 to March 2004.

5.3. Neural Nets

Feed-forward neural networks provide another nonlinear generalization of the autoregression model that has been demonstrated to work well in suitable applications (Faraway and Chatfield, 1998; Hornik and Leisch, 2001; Kajitani et al., 2005). Modeling and forecasting are easily done using **met** (Ripley, 2011). A feed-forward neural net that generalizes the linear autoregressive model of order p may be written,

$$y_t = f_o \left(a + \sum_{i=1}^p \Omega_i x_i + \sum_{j=1}^H w_j f \left(\alpha_j + \sum_{i=1}^p \omega_{i,j} x_{t-i} \right) \right), \quad (6)$$

where \hat{y}_t is the predicted time series at time t and y_{t-1}, \dots, y_{t-p} are the lagged inputs, f_o is the activation function for the output node, f is the activation function for each of the H hidden nodes, $\omega_{i,j}$ are the p weights along the connection for the j -th hidden node, Ω_i is the weight in the skip-layer connection, and a is the bias connection. There are $m(1 + H(p + 2))$ unknown parameters that must be estimated. The hyperparameter H , the number of hidden nodes, is determined by a type of cross-validation and is discussed by Faraway and Chatfield (1998); Hornik and Leisch (2001); Kajitani et al. (2005) in the time series context. The activation functions f and f_o are often chosen to be logistic, $\ell(x) = 1/(1 + e^{-x})$. A schematic illustration for $p = 2$ and $H = 2$ is shown in Figure 20. Feed-forward neural nets may be generalized for multivariate time series.

Hastie et al. (2009) pointed out that the feed-forward neural net defined in eqn. (6) is mathematically equivalent to the projection pursuit regression model. The net defined in eqn. (6) as well as the one illustrated in Figure 20 has just one hidden layer with p and $p = 2$ nodes, respectively. These nets may be generalized to accommodate more than one hidden layer and such nets provide additional flexibility. Ripley (1996) shows that asymptotically for a suitable number of hidden nodes, H , and a large enough training sample, the feed-forward neural net with one hidden layer can approximate any continuous mapping between the inputs and outputs.

6. Unit-root tests

Financial and economic time series such as macro/micro series, stock prices, interest rates and many more, often exhibit nonstationary wandering




Figure 20: A nonlinear version of the AR(2) using the feedforward neural net. This neural net has one hidden layer that is comprised of two hidden nodes. All input nodes have skip-layer connections that connect the input directly with the output.

behavior. Often this type of nonstationarity is easily corrected by differencing and the series is said to have a unit root. Such series are sometimes called homogeneous nonstationary or difference-stationary. Pretesting for a unit root is useful in ARIMA modeling and in cointegration modeling. Since actual time series may also exhibit other departures from the stationary Gaussian ARMA, many other types of unit-root tests have been developed that are appropriate under various other assumptions (Said and Dickey, 1984; Phillips and Perron, 1988; Elliott et al., 1996; Kwiatkowski et al., 1992). State-of-the-art testing for unit roots requires a full model building approach that includes taking into account not only possible general autocorrelation effects but also stochastic and deterministic drift components. An incorrect conclusion may be reached if these effects are not taken into account. Such state-of-the-art tests are implemented in the R packages **fUnitRoots** (Wuertz et al., 2009b) and **urca** (Pfaff, 2010a).

6.1. Overview of the **urca** package

The **urca** (Pfaff, 2010a) package offers a comprehensive and unified approach to unit root testing that is fully discussed in the book Pfaff (2006). The textbook by Enders (2010) also provides an excellent overview

of the state-of-the-art in unit root testing. A useful flowchart for using the **urca** package to test for unit roots is given by Pfaff (2006, Chapter 5).

Three regressions with autocorrelated AR(p) errors are considered for the unit root problem,

$$\Delta Z_t = \beta_0 + \beta_1 t + \gamma Z_{t-1} + \sum_{i=1}^{p-1} \delta_i \Delta Z_{t-i} + e_t \quad (7)$$

$$\Delta Z_t = \beta_0 + \gamma Z_{t-1} + \sum_{i=1}^{p-1} \delta_i \Delta Z_{t-i} + e_t, \quad (8)$$

$$\Delta Z_t = \gamma Z_{t-1} + \sum_{i=1}^{p-1} \delta_i \Delta Z_{t-i} + e_t, \quad (9)$$

corresponding respectively to a unit root:

1. with drift term plus deterministic trend,
2. random walk with drift,
3. pure random walk.

The test for unit root corresponds to an upper-tail test of $\mathcal{H}_0 : \gamma = 0$. The parameters β_0 and β_1 correspond to the drift constant and the deterministic time trend respectively. When $p = 1$, the test reduces to the standard Dickey-Fuller test. To perform the unit-root test, the correct model needs to be identified and the parameters need to be estimated.

The order of the autoregression is estimated using the AIC or BIC. For all three models, the unit-root test is equivalent to testing $\mathcal{H}_0 : \gamma = 0$ is

$$\tau_i = \frac{\hat{\phi} - 1}{\text{SE}(\hat{\phi})}, \quad i = 1, 2, 3,$$

where i denotes the model (9), (8), or (7) respectively. The distribution of τ_i has been obtained by Monte-Carlo simulation or by response surface regression methods (MacKinnon, 1996).

If τ_3 is insignificant, so that $\mathcal{H}_0 : \gamma = 0$ is not rejected, the nonstandard F -statistics Φ_3 and Φ_2 are evaluated using the extra-sum-of-squares principle to test the null hypotheses $\mathcal{H}_0 : (\beta_0, \beta_1, \gamma) = (\beta_0, 0, 0)$ and

$\mathcal{H}_0 : (\beta_0, \beta_1, \gamma) = (0, 0, 0)$ respectively. That is, to test whether the deterministic time trend term is needed in the regression model (eqn 7).

If τ_2 is insignificant, so that $\mathcal{H}_0 : \gamma = 0$ is not rejected, the nonstandard F -statistic Φ_1 is evaluated using the extra-sum-of-squares principle to test the hypotheses $\mathcal{H}_0 : (\beta_0, \gamma) = (0, 0)$. That is, to test whether the regression model has a drift term.

If $\mathcal{H}_0 : \gamma = 0$ is not rejected in the final selected model, we conclude that the series has a unit root.

These steps may be repeated after differencing the series to test if further differencing is needed.

6.1.1. Illustrative example

As an example, consider the U.S. real GNP from 1909 to 1970 in billions of U.S. dollars. From Figure 21, we see the strong upward trend. Since the trend does not appear to follow a straight line, a difference-stationary time series model is suggested. This data set is available as `nporg` in the `urca`




Figure 21: Real U.S. GNP for 1909-1970.

package. We set the maximum lag to 4 and use the BIC to select the optimum number of lags. The code snippet is shown below,

```
R >require("urca")
R >data(nporg)
R >gnp <- na.omit(nporg[, "gnp.r"])
R >summary(ur.df(y = gnp, lags = 4, type = "trend",
+       selectlags = "BIC"))
```

```
#####
# Augmented Dickey-Fuller Test Unit Root Test #
#####
```

Test regression trend

```
Call:
lm(formula = z.diff ~ z.lag.1 + 1 + tt + z.diff.lag)
```

```
Residuals:
    Min       1Q   Median       3Q      Max
-47.149  -9.212   0.819  11.031  23.924
```

```
Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept) -1.89983     4.55369  -0.417  0.67821
z.lag.1      -0.05322     0.03592  -1.481  0.14441
tt           0.74962     0.36373   2.061  0.04423 *
z.diff.lag   0.39082     0.13449   2.906  0.00533 **
```

```
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 15.19 on 53 degrees of freedom
Multiple R-squared: 0.2727,      Adjusted R-squared: 0.2316
F-statistic: 6.625 on 3 and 53 DF,  p-value: 0.0006958
```

Value of test-statistic is: -1.4814 3.8049 2.7942

```
Critical values for test statistics:
    1pct  5pct 10pct
```

```
tau3 -4.04 -3.45 -3.15
phi2  6.50  4.88  4.16
phi3  8.73  6.49  5.47
```

The above R script fit the full model in eqn. (7) with $p = 4$ and used the BIC to select the final model with $p = 1$. Notice that all test statistics are displayed using the `summary` method.

```
#####
# Augmented Dickey-Fuller Test Unit Root Test #
#####
```

Test regression trend

Call:

```
lm(formula = z.diff ~ z.lag.1 + 1 + tt + z.diff.lag)
```

Residuals:

Min	1Q	Median	3Q	Max
-47.374	-8.963	1.783	10.810	22.794

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-0.33082	4.02521	-0.082	0.93479
z.lag.1	-0.04319	0.03302	-1.308	0.19623
tt	0.61691	0.31739	1.944	0.05697 .
z.diff.lag	0.39020	0.13173	2.962	0.00448 **

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 14.88 on 56 degrees of freedom

Multiple R-squared: 0.2684, Adjusted R-squared: 0.2292

F-statistic: 6.847 on 3 and 56 DF, p-value: 0.0005192

Value of test-statistic is: -1.308 3.7538 2.6755

Critical values for test statistics:

	1pct	5pct	10pct
tau3	-4.04	-3.45	-3.15
phi2	6.50	4.88	4.16
phi3	8.73	6.49	5.47

When Sweave (Leisch, 2002) is used, Table 2 may be obtained directly from the output produced in R. Figure 22 shows the graphical model diagnostics.

Table 2: Regression with constant and trend for the U.S. real GNP data starting at 1909 until 1970.

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-0.331	4.025	-0.082	0.935
z.lag.1	-0.043	0.033	-1.308	0.196
tt	0.617	0.317	1.944	0.057
z.diff.lag	0.390	0.132	2.962	0.004




Figure 22: Residual diagnostic of U.S. real GNP data from 1909 to 1970.

The τ_3 statistic for the null hypothesis $\gamma = 0$ is -1.308 and its corresponding critical values at levels 1%, 5%, and 10% with 62 observations are given in Table 3 as -4.04 , -3.45 , and -3.15 respectively. At these levels we can't reject the null hypothesis that $\gamma = 0$ and so we conclude that there is a unit root. Instead of comparing the test statistic value with the critical

Table 3: Critical values for test statistics for drift and trend case eqn. (efADFtest1).

	1pct	5pct	10pct
tau3	-4.04	-3.45	-3.15
phi2	6.50	4.88	4.16
phi3	8.73	6.49	5.47

ones, one can use the MacKinnon's p-value determined from response surface regression methodology (MacKinnon, 1996). The function `punitroot()` is available in `urca`. In the present example, the p-value is 0.88 and it corresponds to the τ_3 statistic value confirming that the unit root hypothesis cannot be rejected as in the code snippet below,

```
R >punitroot(result1.ADF@teststat[1], N = length(gnp),
+ trend = "ct", statistic = "t")
```

```
[1] 0.8767738
```

The F -statistic Φ_3 is used to test whether the deterministic time trend term is needed in the regression model provided that the model has a drift term. The test statistic has a value of 2.68. From Table 3, the critical values of Φ_3 at levels 1%, 5%, and 10% with 62 observations are 8.73, 6.49, and 5.47. We conclude that the null hypothesis is not rejected and a trend term is not needed. Thus we proceed to the next step and estimate the regression parameters in eqn. (8) with a drift term.

```
#####
# Augmented Dickey-Fuller Test Unit Root Test #
#####
```

```
Test regression drift
```

```
Call:
lm(formula = z.diff ~ z.lag.1 + 1 + z.diff.lag)
```

```
Residuals:
```

Min	1Q	Median	3Q	Max
-47.468	-9.719	0.235	10.587	25.192

```
Coefficients:
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.42944	4.01643	0.356	0.7232
z.lag.1	0.01600	0.01307	1.225	0.2257
z.diff.lag	0.36819	0.13440	2.739	0.0082 **

```
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

```
Residual standard error: 15.24 on 57 degrees of freedom
Multiple R-squared: 0.219, Adjusted R-squared: 0.1916
F-statistic: 7.993 on 2 and 57 DF, p-value: 0.0008714
```

```
Value of test-statistic is: 1.2247 3.5679
```

```
Critical values for test statistics:
```

	1pct	5pct	10pct
tau2	-3.51	-2.89	-2.58
phi1	6.70	4.71	3.86

The τ_2 statistic for the null hypothesis $\gamma = 0$ is 1.22474 and its corresponding critical values at levels 1%, 5%, and 10% are given in Table 5 as -3.51 , -2.89 , and -2.58 respectively. From this analysis we conclude that the series behaves like a random walk with a drift constant term. The next question is whether further differencing might be needed. So we simply repeat the unit root modeling and testing using the differenced series as input.

The τ_3 statistic equals to -4.35 . From Table 6, we reject the null hypothesis at 1% and assume that no further differencing is needed.

Table 4: Regression with drift constant for the U.S. real GNP data.

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.42944	4.01643	0.35590	0.72323
z.lag.1	0.01600	0.01307	1.22474	0.22571
z.diff.lag	0.36819	0.13440	2.73943	0.00820

Table 5: Dickey-Fuller critical values for test statistics with drift case.

	1pct	5pct	10pct
tau2	-3.51	-2.89	-2.58
phi1	6.70	4.71	3.86

6.2. Covariate augmented tests

The **CADFtest** package (Lupi, 2011) implements Hansen's covariate augmented Dickey-Fuller test (Hansen, 1995) by including stationary covariates in the model equations,

$$a(L)\Delta Z_t = \beta_0 + \beta_1 t + \gamma Z_{t-1} + b(L)' \Delta X_t + e_t \quad (10)$$

$$a(L)\Delta Z_t = \beta_0 + \gamma Z_{t-1} + b(L)' \Delta X_t + e_t, \quad (11)$$

$$a(L)\Delta Z_t = \gamma Z_{t-1} + b(L)' \Delta X_t + e_t. \quad (12)$$

where $a(L) = 1 - a_1 L + \dots + a_p L^p$ and $b(L)' = b_{q_2} L^{-q_2} + \dots + b_{q_1} L^{q_1}$. If the main function `CADFtest()` is applied without any stationary covariates, the ordinary ADF test is performed. In the illustrative example below, taken from the `CADFtest()` online documentation, the augmented test strongly rejects the unit root hypothesis, with a p-value less than 2%. On the other hand, with the covariate, the test produces a p-value of about 9%. This is shown in the the R session below,

```
R >require(CADFtest)
R >data(npext, package = "urca")
R >npext$unemrate <- exp(npext$unemploy)
R >L <- ts(npext, start = 1860)
R >D <- diff(L)
R >S <- window(ts.intersect(L, D), start = 1909)
```


Table 6: Critical values for test statistics testing for second differences.

	1pct	5pct	10pct
tau3	-4.04	-3.45	-3.15
phi2	6.50	4.88	4.16
phi3	8.73	6.49	5.47

```
R >CADFtest(L.gnpperca ~ D.unemrate, data = S, max.lag.y = 3,
+ kernel = "Parzen", prewhite = FALSE)
```

CADF test

```
data: L.gnpperca ~ D.unemrate
CADF(3,0,0) = -3.413, rho2 = 0.064, p-value =
0.001729
alternative hypothesis: true delta is less than 0
sample estimates:
delta
-0.08720302
```

7. Cointegration and VAR models

In the simplest case, two time series that are both difference-stationary are said to be cointegrated when a linear combination of them is stationary. Some classic examples ([Engle and Granger, 1987](#)) of bivariate cointegrated series include:

- consumption and income
- wages and prices
- short and long term interest rates

Further examples are given in most time series textbooks with an emphasis on economic or financial series ([Enders, 2010](#); [Chan, 2010](#); [Tsay, 2010](#); [Lütkepohl, 2005](#); [Hamilton, 1994](#); [Banerjee et al., 1993](#)).

A cointegration analysis requires careful use of the methods discussed in these books since spurious relationships can easily be found when working with difference-stationary series (Granger and Newbold, 1974). Most financial and economic time series are not cointegrated. Cointegration implies a deep relationship between the series that is often of theoretical interest in economics. When a cointegrating relationship exists between two series, Granger causality must exist as well (Pfaff, 2006). The **vars** package (Pfaff, 2010b) for vector autoregressive modeling is described in the book (Pfaff, 2006) and article (Pfaff, 2008). This package, along with its companion package **urca** (Pfaff, 2010a), provides state-of-the-art methods for cointegration analysis and modeling stationary and nonstationary multivariate time series.

Full support for modeling, forecasting and analysis tools are provided for the vector autoregressive time series model (VAR), structural VAR (SVAR) and structural vector error-correction models (SVEC). The VAR(p) stationary model for a k -dimensional time series, $\{\mathbf{y}_t\}$

$$\mathbf{y}_t = \boldsymbol{\delta}\mathbf{d}_t + \boldsymbol{\Phi}_1\mathbf{y}_{t-1} + \dots + \boldsymbol{\Phi}_p\mathbf{y}_{t-p} + \boldsymbol{\epsilon}_t, \quad (13)$$

where $\boldsymbol{\delta}, \boldsymbol{\Phi}_\ell = (\phi_{ij,\ell})_{k \times k}$ are coefficient matrices, \mathbf{d}_t is a matrix containing a constant term, linear trend, seasonal indicators or exogenous variables, and $\boldsymbol{\epsilon}_t \sim N(\mathbf{0}, I_k)$. Using the **vars** package, the VAR model is estimated using OLS. The basic VAR model, without the covariates \mathbf{d}_t , may also be estimated using the R core function `ar()`. In the case of the SVAR model,

$$\mathbf{A}\mathbf{y}_t = \boldsymbol{\delta}\mathbf{d}_t + \boldsymbol{\Phi}_1\mathbf{y}_{t-1} + \dots + \boldsymbol{\Phi}_p\mathbf{y}_{t-p} + \mathbf{B}\boldsymbol{\epsilon}_t, \quad (14)$$

where \mathbf{A} , and \mathbf{B} are $k \times k$ matrices. With the structural models, further restrictions are needed on the parameters and after the model has been uniquely specified, it is estimated by maximum likelihood. The SVEC model is useful for modeling non-stationary multivariate time series and is an essential tool in cointegration analysis. The basic error correction model, VEC, may be written,

$$\nabla\mathbf{y}_t = \boldsymbol{\Pi}\mathbf{y}_t + \boldsymbol{\Gamma}_1\nabla\mathbf{y}_{t-1} + \dots + \nabla\boldsymbol{\Gamma}_p\mathbf{y}_{t-p+1} + \boldsymbol{\epsilon}_t, \quad (15)$$

where ∇ is the first-differencing operator and $\boldsymbol{\Pi}$ and $\boldsymbol{\Gamma}_\ell, \ell = 1, \dots, p-1$ are parameters. As with the VAR model, the VEC model may be generalized

to the SVEC model with coefficient matrices \mathbf{A} and/or \mathbf{B} . A cointegration relationship exists provided that $0 < \text{rank } \mathbf{\Pi} < p$. When $\text{rank } \mathbf{\Pi} = 0$, a VAR model with the first differences may be used and when $\mathbf{\Pi}$ is of full rank, a stationary VAR model of order p is appropriate. The **vars** package includes functions for model fitting, model selection and diagnostic checking as well as forecasting with VAR, SVAR and SVEC models. Cointegration tests and analysis are provided in the **urca**. In addition to the two-step method of [Engle and Granger \(1987\)](#), tests based on the method of [Phillips and Ouliaris \(1990\)](#) and the likelihood method ([Johansen, 1995](#)) are implemented in the **urca** package. Illustrative examples of how to use the software for multivariate modeling and cointegration analysis are discussed in the book, paper and packages of [Pfaff \(2006, 2008, 2010b\)](#).

8. GARCH time series

Volatility refers to the random and autocorrelated changes in variance exhibited by many financial time series. The GARCH family of models ([Engle, 1982](#); [Bollerslev, 1986](#)) capture quite well volatility clustering as well as the thick-tailed distributions often found with financial time series such as stock returns and foreign exchange rates. The GARCH family of models is discussed in more detail in textbooks dealing with financial time series ([Enders, 2010](#); [Chan, 2010](#); [Tsay, 2010](#); [Cryer and Chan, 2008](#); [Shumway and Stoffer, 2011](#); [Hamilton, 1994](#)).

A GARCH(p, q) sequence $a_t, t = \dots, -1, 0, 1, \dots$ is of the form

$$a_t = \sigma_t \epsilon_t$$

and

$$\sigma_t^2 = \alpha_0 + \sum_{i=1}^p \alpha_i a_{t-i}^2 + \sum_{j=1}^q \beta_j \sigma_{t-j}^2,$$

where $\alpha_0 > 0$, $\alpha_i \geq 0$, $1 \leq i \leq p$, $\beta_j \geq 0$, $1 \leq j \leq q$ are parameters. The errors ϵ_t are assumed to be independent and identically distributed from a parametric distribution such as normal, generalized error distribution (GED), Student-t or skewed variations of these distributions. While ARMA models deal with nonconstant conditional expectation, GARCH models handle non-constant conditional variance. Sometimes those two models are combined to form the ARMA/GARCH family of models. A comprehensive

account of these models is also given in the book by [Zivot and Wang \(2006\)](#). This book also serves as the documentation for the well-known S-Plus add-on module, **Finmetrics**. Many of the methods provided by **Finmetrics** for GARCH and related models are now available with the **fGARCH** package ([Wuertz et al., 2009a](#)). In the following, we give a brief discussion of the use of **fGARCH** for simulation, fitting and inferences. The principal functions in this package include `garchSpec`, `garchSim`, and `garchFit` and related methods functions. The **fGarch** package allows for a variety of distributional assumptions for the error sequence ϵ_t . As an illustrative example, we simulate a GARCH(1,1) with $\alpha_0 = 10^{-6}$, $\alpha_1 = 0.2$, and $\beta_1 = 0.7$ and with a skewed GED distribution with skewness coefficient 1.25 and shape parameter 4.8. The simulated series is shown in [Figure 23](#).

```
R> require("fGarch")
R> spec <- garchSpec(model = list(omega = 1e-06, alpha = 0.2,
+   beta = 0.7, skew = 1.25, shape = 4.8), cond.dist = "sged")
R> x <- garchSim(spec, n = 1000)
```




Figure 23: Simulated GARCH(1,1) with $\alpha_0 = 10^{-6}$, $\alpha_1 = 0.2$, $\beta_1 = 0.7$.

To fit the above simulated data with GARCH(1,1) we could use,

```
R> out <- garchFit(~garch(1, 1), data = x, trace = FALSE)
```

Some of the inferences that can be carried out by using the `summary()` function, include the Jarque-Bera and Shapiro-Wilk normality tests, various Ljung-Box white noise tests, and ARCH effect tests.

As a further illustration, we fit an ARMA/GARCH model to the U.S. inflation (Bollerslev, 1986). We used the GNP deflator for 1947-01-01 to 2010-04-01. There were $n = 254$ observations which are denoted by $z_t, t = 1, \dots, n$. Then the inflation rate may be estimated by the logarithmic difference, $r_t = \log(z_t) - \log(z_{t-1})$. The following ARMA/GARCH model was fit using the function `garchFit()` in **fGarch**,
 $r_t = 0.103 + 0.369r_{t-1} + 0.223r_{t-2} + 0.248r_{t-3} + \epsilon_t$, and
 $\sigma_t^2 = 0.004 + 0.269\epsilon_{t-1}^2 + 0.716\sigma_{t-1}^2$. Figure 24 shows time series plots for r_t and σ_t . The `tseries` (Trapletti, 2011) can also fit GARCH models but




Figure 24: Inflation rate, r_t , and volatility, σ_t .

fGarch provides a more comprehensive approach.

9. Wavelet methods in time series analysis

Consider a time series of dyadic length, $z_t, t = 1, \dots, n$, where $n = 2^J$. The discrete wavelet transformation (DWT) decomposes the time series into J wavelet coefficients vectors, $W_j, j = 0, \dots, J - 1$ each of length

$n_j = 2^{J-j}$, $j = 1, \dots, J$ plus a scaling coefficient V_j . Each wavelet coefficient is constructed as a difference of two weighted averages each of length $\lambda_j = 2^{j-1}$. Like the discrete Fourier transformation, the DWT provides an orthonormal decomposition, $W = \mathcal{W}Z$, where $W' = (W'_1, \dots, W'_{J-1}, V'_{J-1})$, $Z = (z_1, \dots, z_n)'$ and \mathcal{W} is an orthonormal matrix. In practice, the DWT is not computed using matrix multiplication but much more efficiently using filtering and downsampling (Percival and Walden, 2000, Ch 4). The resulting algorithm is known as the pyramid algorithm and computationally it is even more efficient than the fast Fourier transform. Applying the operations in reverse order yields the inverse DWT. Sometimes a partial transformation is done, producing the wavelet coefficient vectors W_j , $j = 0, \dots, J_0$, where $J_0 < J - 1$. In this case, the scaling coefficients are in the vector, V_{J_0} of length 2^{J-J_0} . The wavelet coefficients are associated with changes in the time series over the scale $\lambda_j = 2^{j-1}$ while the scaling coefficients, V_{J_0} , are associated with the average level on scale $\tau = 2^{J_0}$. The maximum overlap DWT or MODWT omits the downsampling. The MODWT has many advantages over the DWT (Percival and Walden, 2000, Ch 5) even though it does not provide an orthogonal decomposition. Percival and Walden (2000) provide an extensive treatment of wavelet methods for time series research with many interesting scientific time series. Gençay et al. (2002) follows a similar approach to wavelets as given by Percival and Walden (2000) but with an emphasis on financial and economic applications.

All important methods as well as all datasets discussed in the books by Percival and Walden (2000); Gençay et al. (2002) are available in the R packages `waveslim` (Whitcher, 2010) and `wmtsa` (Constantine and Percival, 2010). Nason (2008) provides a general introduction to wavelet methods in statistics, including smoothing and multiscale time series analysis. R scripts are used extensively in his book and all figures in the book (Nason, 2008) may be reproduced using R scripts available in the `wavethresh` R package (Nason, 2010).

Figure 25 shows the denoised annual Nile riverflows (Hipel and McLeod, 1994) using the universal threshold with hard thresholding and Haar wavelets. Hipel and McLeod (1994); Hipel et al. (1975) fit a step intervention analysis time series model with AR(1) noise. Physical reasons as well as cumsum analysis were presented (Hipel and McLeod, 1994, §19.2.4) to suggest 1903 as the start of intervention that was due to the operation of the Aswan dam. The fitted step intervention is represented by

the three line segments while the denoised flows are represented by the jagged curve. The points show actual observed flows. Figure 25 suggests the intervention actually may have started a few years prior to 1903. The computations for Figure 25 were done using the functions `modwt()`, `universal.thresh.modwt()` and `imodwt()` in the package `waveslim`.




Figure 25: Mean annual Nile flow, October to September, Aswan.

An estimate of the wavelet variance, $\hat{\sigma}^2(\lambda_j)$ is obtained based on the variance of the wavelet coefficients in an MODWT transformation at scale $\lambda_j = 2^{j-1}$. The wavelet variance is closely related to the power spectral density function and

$$\hat{\sigma}^2(\lambda_j) \approx 2 \int_{1/\lambda_j}^{2/\lambda_j} p(f)df.$$

The wavelet variance decomposition for the annual sunspot numbers, `sunspot.year` in R is shown in Figure 26. This figure was produced using the `wavVar` function in `wmtsa` and the associated plot method. The 95%

confidence intervals are shown in Figure 26. The wavelet variances correspond to changes over 1, 2, 4, 8 and 16 years.




Figure 26: Wavelet variance, yearly sunspot numbers, 1700-1988.

Multiresolution analysis (MRA) is another widely useful wavelet method for time series analysis. The MRA decomposition works best with the MODWT. The `mra` function in `waveslim` was used to produce the decomposition of an electrocardiogram time series that is shown in Figure 27. The `1a8` or least-asymmetric filter with half-length 8 was used (Percival and Walden, 2000, p. 109). A similar plot is given by Percival and Walden (2000, Figure 184).




Figure 27: MRA using MODWT with 1a8 filter. ECG time series comprised of about 15 beats of a human heart, sampled at 180 Hz, units are millivolts and $n = 2048$.

10. Stochastic differential equations (SDE)

A SDE is comprised of a differential equation that includes a stochastic process, the simplest example being Brownian motion. Geometrical Brownian motion is often used to describe stock market prices. This SDE may be written, $dP(t) = P(t)\mu dt + P(t)\sigma dW(t)$ where $P(t)$ is the price at time t and the parameters $\mu > 0$ and $\sigma > 0$ are the drift and diffusion parameters. The Gaussian white noise term, $W(t)$, may be considered the derivative of Brownian motion. This SDE may also be written, $d\log(P(t)) = \mu dt + \sigma dW(t)$, so we see that $P(t) > 0$ and $\log(P(t))$ is Brownian motion.

More complicated SDE's may involve more complex drift and volatility functions. The book (Iacus, 2008) provides an intuitive and informal introduction to SDE and could be used in an introductory course on SDE. Only SDE's with Gaussian white noise are considered. The accompanying R package (Iacus, 2009) provides R scripts for all figures in the book (Iacus, 2008) as well as functions for simulation and statistical inference with SDE.

An important area of application is in financial mathematics where option values or risk assessments are often driven by SDE systems. Usually Monte Carlo simulation is the only way to find approximate solutions. The main class of SDE considered by this package is a diffusion process of the following form,

$$dX(t) = b(t, X(t)) dt + \sigma(t, X(t)) dW(t) \quad (16)$$

with some initial condition $X(0)$, where $W(t)$ is a standard Brownian motion. According to Itô formula, (16) can be represented as

$$X(t) = X(0) + \int_0^t b(u, X(u)) du + \int_0^t \sigma(u, X(u)) dW(u).$$

Under some regular conditions on the drift $b(\cdot, \cdot)$ and diffusion $\sigma^2(\cdot, \cdot)$, (16) has either a unique strong or weak solution. In practice, the class of SDE given by (16) is too large. The following diffusion process covers many well-known and widely used stochastic processes, including Vasicek (VAS), Ornstein-Uhlenbeck (OU), Black-Scholes-Merton (BS) or geometric Brownian motion, and Cox-Ingersoll-Ross (CIR),

$$dP(t) = P(t)\mu dt + P(t)\sigma dW(t) \quad dX(t) = b(X(t)) dt + \sigma(X(t)) dW(t). \quad (17)$$

The main function is `sde.sim()` and it has extensive options for the general diffusion process (17) or more specific processes. The function `DBridge()` provides another general purpose function for simulating diffusion bridges. Simple to use functions for simulating a Brownian bridge and geometric Brownian motion, `BBridge()` and `GBM()`, are also provided. Using `sde.sim()`, we simulate ten replications of Brownian motions each starting at the $X(0) = 0$ and comprised of 1000 steps. The results are displayed in Figure 28.




Figure 28: Ten Brownian motions.

A more complex SDE,

$$dX(t) = (5 - 11x + 6x^2 - x^3)dt + dW(t)$$

with $X(0) = 5$ is simulated using three different algorithms and using two different step-sizes $\Delta = 0.1$ and $\Delta = 0.25$. For the smaller step size $\Delta = 0.1$, Figure 29 suggests all three algorithms work about equally well. But only the Shoji-Ozaki algorithm appears to work with the larger step size $\Delta = 0.25$.

In addition to simulation, the `sde` package provides functions for parametric and nonparametric estimation: `EULERloglik()`, `ksmooth()`,




Figure 29: Simulations of $dX(t) = (5 - 11x + 6x^2 - x^3)dt + dW(t)$ using three different algorithms and two different step sizes.

`SIMloglik()`, and `simple.ef()`. Approximation of conditional density $X(t)|X(t_0) = x_0$ at point x_0 of a diffusion process is available with the functions: `dcEulerian()`, `dcEuler()`, `dcKessler()`, `dcozaki()`, `dcShoji()`, and `dcSim()`.

11. Conclusion

There are many more packages available for time series than discussed in this article and many of these are briefly described in the CRAN Task Views.¹⁰ In particular, see task views for **Econometrics**, **Finance** and **TimeSeries**. We have selected those packages that might be of most general interest, that have been most widely used and that we are most familiar with. The reader should note that the packages published on CRAN, including those in the task views, need only obey formatting rules and not produce computer errors. There is no endorsement that packages available on CRAN produce correct or useful results. On the other hand,

¹⁰<http://cran.r-project.org/web/views/>

packages discussed in the *Journal of Statistical Software* or published by major publishers such as Springer-Verlag or Chapman & Hall/CRC have been carefully reviewed for correctness and quality.

Researchers wishing to increase the impact of their work should consider implementing their methods in R and making it available as a package on CRAN. Developing R packages is discussed in the online publication by [R Development Core Team \(2011\)](#) and from a broader perspective by [Chambers \(2008\)](#).

Acknowledgements

Drs. A. I. McLeod and Hao Yu would like to thank NSERC for Discovery Grants awarded to each of us. The authors would also like to thank Achim Zeileis for some suggestions and an anonymous referee for their comments.

12. Appendix

12.1. datasets

Dataset name	Description
AirPassengers	monthly airline passengers, 1949-1960
BJsales	sales data with leading indicator
BOD	biochemical oxygen demand
EuStockMarkets	daily close price, European stocks, 1991-1998
LakeHuron	level of Lake Huron 1875-1972
Nile	flow of the river Nile
UKDriverDeaths	road casualties, Great Britain 1969-84
UKgas	UK quarterly gas consumption
USAccDeaths	accidental deaths in the US 1973-1978
USPersonalExpenditure	personal expenditure data
WWWusage	internet usage per minute
WorldPhones	the world's telephones
airmiles	passenger miles, US airlines, 1937-1960
austres	quarterly time series, Australian residents
co2	mauna loa atmospheric co2 concentration
UKLungDeaths	monthly deaths from lung diseases in the UK
freeny	Freeny's revenue data
longley	Longley's economic regression data
lynx	annual Canadian lynx trappings 1821-1934
nhtemp	average yearly temperatures in New Haven
nottem	monthly temperature, Nottingham, 1920-39
sunspot.month	monthly sunspot data, 1749-1997
sunspot.year	yearly sunspot data, 1700-1988
sunspots	monthly sunspot numbers, 1749-1983
treering	yearly treering data, -6000-1979
uspop	populations recorded by the US census

Table 7: **datasets** time series data.

12.2. stats

Function	Purpose
<code>embed</code>	matrix containing lagged values
<code>lag</code>	lagged values
<code>ts</code>	create a time series object
<code>ts.intersect</code>	intersection, multivariate series by
<code>ts.union</code>	union, multivariate series by union
<code>time</code>	extract time from a ts-object
<code>cycle</code>	extract seasonal times from a ts-object
<code>frequency</code>	sampling interval
<code>window</code>	select subset of time series

Table 8: **stats** utilities for ts-objects. These functions are useful for creating and manipulating univariate and multivariate time series.

Function	Purpose
<code>acf</code>	acf, pacf
<code>ccf</code>	cross-correlation
<code>cpgram</code>	Bartlett's cumulate periodogram test
<code>lag.plot</code>	alternative time series plot
<code>fft</code>	fast Fourier transform
<code>convolve</code>	convolution via fft
<code>filter</code>	moving-average/autoregressive filtering
<code>spectrum</code>	spectral density estimation
<code>toeplitz</code>	Toeplitz matrix

Table 9: **stats** autocorrelation and spectral analysis functions.

Function	Purpose
<code>arima</code> , <code>arima0</code>	fit ARIMA
<code>ar</code>	fit AR
<code>KalmanLike</code>	loglikelihood, univariate state-space model
<code>KalmanRun</code>	KF filtering
<code>KalmanSmooth</code>	KF smoothing
<code>KalmanForecast</code>	KF forecasting
<code>makeARIMA</code>	ARIMA to KF
<code>PP.test</code>	Phillips-Perron unit root test
<code>tsdiag</code>	diagnostic checks
<code>ARMAacf</code>	theoretical ACF of ARMA
<code>acf2AR</code>	fit AR to ACF
<code>Box.test</code>	Box-Pierce or Ljung-Box test
<code>diff</code> , <code>diffinv</code>	difference or inverse
<code>ARMAtoMA</code>	MA expansion for ARMA
<code>arima.sim</code>	simulate ARIMA
<code>HoltWinters</code>	Holt-Winters filtering
<code>StructTS</code>	Kalman filter modeling

Table 10: **stats** functions for time series models. In addition many of these function have `predict` and `residuals` methods.

Function	Purpose
<code>filter</code>	moving-average/autoregressive filtering
<code>tsSmooth</code>	smooth from <code>StructTS</code> object
<code>stl</code>	seasonal-trend-loess decomposition
<code>decompose</code>	seasonal decomposition, moving-average filters

Table 11: **stats** smoothing and filtering.

12.3. `tseries`

Function	Purpose
<code>adf.test</code>	augmented Dickey-Fuller test
<code>bds.test</code>	Breusch-Godfrey test
<code>garch</code>	fit GARCH models to time series
<code>get.hist.quote</code>	download historical finance data
<code>jarque.bera.test</code>	Jarque-Bera test
<code>kpss.test</code> KPSS	test for stationarity
<code>quadmap</code>	quadratic map (logistic equation)
<code>runs.test</code>	runs test
<code>terasvirta.test</code>	Teraesvirta neural network test for nonlinearity
<code>tsbootstrap</code>	bootstrap for general stationary data
<code>white.test</code>	White neural network test for nonlinearity

Table 12: `tseries` functions.

Dataset name	Description
<code>bev</code>	Beveridge wheat price index, 1500-1869
<code>camp</code>	Mount Campito, treering data, -3435-1969
<code>ice.river</code>	Icelandic river Data
<code>NelPlo</code>	Nelson-Plosser macroeconomic time series
<code>nino</code>	sea surface temperature, El Niño indices
<code>tcm</code>	monthly yields on treasury securities
<code>tcmd</code>	daily yields on treasury securities
<code>USEconomic</code>	U.S. economic variables

Table 13: `tseries` time series data.

12.4. Forecast

Function	Purpose
<code>accuracy()</code>	accuracy measures of forecast
<code>BoxCox, invBoxCox()</code>	Box-Cox transformation
<code>decompose()</code>	improved version of <code>decompose()</code>
<code>dm.test()</code>	Diebold-Mariano test compares the forecast accuracy
<code>forecast()</code>	generic function with various methods
<code>monthdays()</code>	number of days in seasonal series
<code>na.interp()</code>	interpolate missing values
<code>naive(), snaive()</code>	ARIMA(0,1,0) forecast and seasonal version
<code>seasadj()</code>	seasonally adjusted series
<code>seasonaldummy()</code>	create matrix of seasonal indicator variables
<code>seasonplot()</code>	season plot

Table 14: General purpose utility functions.

Function	Purpose
<code>arfima</code>	automatic ARFIMA
<code>Arima</code>	improved version of <code>arima()</code>
<code>arima.errors</code>	removes regression component
<code>auto.arima</code>	automatic ARIMA modeling
<code>ndiffs</code>	use unit root test to determine differencing
<code>tsdisplay()</code>	display with time series plot, ACF, PACF, etc.

Table 15: ARIMA functions.

Function	Purpose
<code>croston</code>	exponential forecasting for intermittent series
<code>ets</code>	exponential smoothing state space model
<code>logLik.ets</code>	loglikelihood for ets object
<code>naive()</code> , <code>snaive()</code>	ARIMA(0,1,0) forecast and seasonal version
<code>rwf()</code>	random walk forecast with possible drifts
<code>ses()</code> , <code>holt()</code> , <code>hw()</code>	exponential forecasting methods
<code>simulate.ets()</code>	simulation method for ets object
<code>sindexf</code>	seasonal index, future periods
<code>splinef</code>	forecast using splines
<code>thetaf</code>	forecast using theta method
<code>tslm()</code>	<code>lm()</code> -like function using trend and seasonal

Table 16: Exponential smoothing and other time series modeling functions.

12.5. **itsa**

Function	Purpose
<code>DHSimulate</code>	simulate using Davies-Harte method
<code>DLLoglikelihood</code>	exact concentrated log-likelihood
<code>DLResiduals</code>	standardized prediction residuals
<code>DLSimulate</code>	simulate using DL recursion
<code>SimGLP</code>	simulate general linear process
<code>TrenchInverse</code>	Toeplitz matrix inverse
<code>ToeplitzInverseUpdate</code>	updates the inverse
<code>TrenchMean</code>	exact MLE for mean
<code>TrenchForecast</code>	exact forecast and variance

Table 17: Main functions in **itsa**.

12.6. FitAR

Function	Purpose
PacfPlot	partial autocorrelation plot
SelectModel	AIC/BIC selection
TimeSeriesPlot	time series plot

Table 18: **FitAR** model selection functions.

Function	Purpose
FitAR	exact mle for AR(p)/subset ARzeta
FitARLS	LS for AR(p)/subset ARphi
GetFitAR	fast exact mle for AR(p)/subset ARzeta
GetFitARLS	fast LS for AR(p) and subset ARphi
GetARMeanMLE	exact mean MLE in AR
AR1Est	exact MLE for mean-zero AR(1)

Table 19: **FitAR** estimation functions.

Function	Purpose
Boot	generic parametric bootstrap
Boot.FitAR	method for FitAR
Boot.ts	method for ts
LjungBox	Ljung-Box portmanteau test
LBQPlot	plot Ljung-Box test results
RacfPlot	residual acf plot
JarqueBeraTest	test for normality

Table 20: **FitAR** diagnostic check functions.

Function	Purpose
AcfPlot	general purpose correlation plotting
ARSdf	AR spectral density via FFT
ARToMA	impulse coefficients
ARToPacf	transform AR to PACF
BackcastResidualsAR	compute residuals using backforecasting
cts	concatenate time series
InformationMatrixAR	Fisher information matrix AR
InformationMatrixARp	Fisher information matrix subset case, ARp
InformationMatrixARz	Fisher information matrix subset case, ARz
InvertibleQ	test if invertible or stationary-casual
PacfDL	compute PACF from ACF using DL recursions
PacfToAR	transform PACF to AR
sdfplot	generic spectral density plot
sdfplot.FitAR	method for class FitAR
sdfplot.Arima	method for class Arima
sdfplot.ar	method for class ar
sdfplot.ts	method for class ts
sdfplot.numeric	method for class numeric
SimulateGaussianAR	simulate Gaussian AR
Readts	input time series
TacvfAR	theoretical autocovariances AR
TacvfMA	theoretical autocovariances MA
VarianceRacfAR	variance of residual acf, AR
VarianceRacfARp	variance of residual acf, subset case, ARp
VarianceRacfARz	variance of residual acf, subset case, ARz

Table 21: **FitAR** miscellaneous functions.

References

- Adler, J., 2009. R in a Nutshell. O'Reilly, Sebastopol, CA.
- Aknouche, A., Bibi, A., 2009. Quasi-maximum likelihood estimation of periodic garch and periodic arma-garch processes. *Journal of Time Series Analysis* 30 (1), 19–46.
- Baillie, R. T., 1996. Long memory processes and fractional integration in econometrics. *Journal of Econometrics* 73 (1), 5 – 59.
- Banerjee, A., Dolado, J. J., Galbraith, J. W., Hendry, D. F., 1993. *Cointegration, Error Correction, and the Econometric Analysis of Non-Stationary Data*. Oxford University Press, Oxford.
- Becker, R. A., Clark, L. A., Lambert, D., 1994. Cave plots: A graphical technique for comparing time series. *Journal of Computational and Graphical Statistics* 3 (3), 277–283.
- Beran, J., 1994. *Statistics for Long Memory Processes*. Chapman & Hall/CRC, Boca Raton.
- Beran, J., Whitcher, B., Maechler, M., 2009. *longmemo: Statistics for Long-Memory Processes*.
URL <http://CRAN.R-project.org/package=longmemo>
- Bloomfield, P., 2000. *Fourier Analysis of Time Series: An Introduction*, 2nd Edition. Wiley, New York.
- Bollerslev, T., 1986. Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics* 31 (3), 307–327.
- Box, G., Jenkins, G. M., Reinsel, G. C., 2008. *Time Series Analysis: Forecasting and Control*, 4th Edition. Hoboken, N.J. : J.Wiley, New York.
- Box, G. E. P., Tiao, G. C., 1975. Intervention analysis with applications to economic and environmental problems. *Journal of the American Statistical Association* 70 (349), 70–79.
- Braun, W. J., Murdoch, D. J., 2008. *A First Course in Statistical Programming with R*. Cambridge University Press, Cambridge.

- Brockwell, P. J., Davis, R. A., 1991. Time Series: Theory and Methods, 2nd Edition. Springer, New York.
- Brown, R. L., Durbin, J., Evans, J. M., 1975. Techniques for testing the constancy of regression relationships over time. *Journal of the Royal Statistical Society B* 37, 149–163.
- Chambers, J. M., June 2008. Software for Data Analysis: Programming with R. Statistics and Computing. Springer-Verlag.
- Chan, K.-S., 2011. TSA: Time Series Analysis. R package version 0.98. URL <http://CRAN.R-project.org/package=TSA>
- Chan, N. H., 2010. Time Series: Applications to Finance with R, 3rd Edition. Wiley, New York.
- Cleveland, W. S., 1993. Visualizing Data. Hobart Press.
- Cleveland, W. S., McGill, M. E., McGill, R., 1988. The shape parameter of a two-variable graph. *Journal of the American Statistical Association* 83 (402), 289–300.
- Constantine, W., Percival, D., 2010. wmtsa: Insightful Wavelet Methods for Time Series Analysis. R package version 1.0-5. URL <http://CRAN.R-project.org/package=wmtsa>
- Cook, D., Swayne, D. F., 2007. Interactive Dynamic Graphics for Data Analysis R. Springer-Verlag, New York.
- Cowpertwait, P. S., Metcalfe, A. V., 2009. Introductory Time Series with R. Springer Science+Business Media, LLC, New York.
- Craigmile, P. F., 2003. Simulating a class of stationary gaussian processes using the davies-harte algorithm, with application to long memory processes. *Journal of Time Series Analysis* 24, 505–511.
- Crawley, M. J., 2007. The R Book. Wiley, New York.
- Cryer, J. D., Chan, K.-S., 2008. Time Series Analysis: With Applications in R, 2nd Edition. Springer Science+Business Media, LLC, New York.

- Dalgaard, P., 2008. *Introductory Statistics with R*. Springer Science+Business Media, LLC, New York.
- Davies, R. B., Harte, D. S., 1987. Tests for hurst effect. *Biometrika* 74, 95–101.
- Dethlefsen, C., Lundbye-Christensen, S., Christensen, A. L., 2009. *sspir: State Space Models in R*.
URL <http://CRAN.R-project.org/package=sspir>
- Diebold, F. X., Rudebusch, G. D., 1989. Long memory and persistence in aggregate output. *Journal of Monetary Economics* 24, 189–209.
- Durbin, J., Koopman, S. J., 2001. *Time Series Analysis by State Space Methods*. Oxford University Press.
- Elliott, G., Rothenberg, T. J., Stock, J. H., 1996. Efficient tests for an autoregressive unit root. *Econometrica* 64 (4), 813–836.
- Enders, W., 2010. *Applied Econometric Time Series*, 3rd Edition. John Wiley and Sons, New York.
- Engle, R. F., 1982. Autoregressive conditional heteroscedasticity with estimates of the variance of united kingdom inflation. *Econometrica* 50 (4), 987–1007.
- Engle, R. F., Granger, C. W. J., 1987. Co-integration and error correction: Representation, estimation, and testing. *Econometrica* 55 (2), 251–276.
- Everitt, B. S., Hothorn, T., 2009. *A Handbook of Statistical Analyses Using R*, 2nd Edition. Chapman and Hall/CRC, Boca Raton.
- Faraway, J., Chatfield, C., 1998. Time series forecasting with neural networks: A comparative study using the airline data. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 47 (2), 231–250.
- Fox, J., 2005. The R commander: A basic-statistics graphical user interface to R. *Journal of Statistical Software* 14 (9), 1–42.
URL <http://www.jstatsoft.org/v14/i09>

- Fraley, C., Leisch, F., Maechler, M., Reisen, V., Lemonte, A., 2009. fracdiff: Fractionally differenced ARIMA aka ARFIMA(p,d,q) models.
URL <http://CRAN.R-project.org/package=fracdiff>
- Friedman, J. H., 1991. Multivariate adaptive regression splines. *The Annals of Statistics* 19 (1), 1–67.
- Gelper, S., Fried, R., Croux, C., 2010. Robust forecasting with exponential and holt-winters smoothing. *Journal of Forecasting* 29 (3), 285–300.
- Gençay, R., Selçuk, F., Whitcher, B., 2002. *An Introduction to Wavelets and Other Filtering Methods in Finance and Economics*. Academic Press, New York.
- Gentleman, R., 2009. *R Programming for Bioinformatics*. Chapman and Hall/CRC.
- Gentleman, R., Ihaka, R., 1996. R: A language for data analysis and graphics. *The Journal of Computational and Graphical Statistics* 5 (2), 491–508.
- Gilbert, P., 1993. State space and arma models: An overview of the equivalence. Bank of Canada Publications. Working Paper 1993-00.
URL <http://www.bankofcanada.ca/1993/03/publications/research/working-paper-199/>
- Gilbert, P., 2011. dse: Dynamic Systems Estimation (time series package).
URL <http://CRAN.R-project.org/package=dse>
- Granger, C. W. J., Newbold, P., 1974. Spurious regressions in econometrics. *Journal of Econometrics* 2, 111–120.
- Granger, C. W. J., Newbold, P., 1976. Forecasting transformed series. *Journal of the Royal Statistical Society. Series B (Methodological)* 38 (2), 189–203.
- Graves, S., 2011. FinTS: Companion to Tsay (2005) Analysis of Financial Time Series. R package version 0.4-4.
URL <http://CRAN.R-project.org/package=FinTS>

- Grolemund, G., Wickham, H., 2011. Dates and times made easy with lubridate. *Journal of Statistical Software* 40 (3), 1–25.
URL <http://www.jstatsoft.org/v40/i03>
- Hamilton, J. D., 1994. *Time Series Analysis*. Princeton University Press, Princeton, NJ.
- Hansen, B. E., 1995. Rethinking the univariate approach to unit root testing: Using covariates to increase power. *Econometric Theory* 11 (5), 1148–1171.
- Harrison, J., West, M., 1997. *Bayesian forecasting and dynamic models*.
- Harvey, A., 1989. *Forecasting, Structural Time Series Models and the Kalman Filter*. Cambridge University Press, Cambridge.
- Haslett, J., Raftery, A. E., 1989. Space-Time Modelling with Long-Memory Dependence: Assessing Ireland’s Wind Power Resource. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 38 (1), 1–50.
- Hastie, T., Tibshirani, R., 2011. mda: Mixture and Flexible Discriminant Analysis. R package version 0.4-2.
URL <http://CRAN.R-project.org/package=mda>
- Hastie, T., Tibshirani, R., Friedman, J. H., 2009. *The Elements of Statistical Learning*, 2nd Edition. Springer-Verlag, New York.
- Heiberger, R. M., Neuwirth, E., 2009. *R Through Excel: A Spreadsheet Interface for Statistics, Data Analysis, and Graphics*. Springer Science+Business Media, LLC, New York.
- Helske, J., 2011. KFAS: Kalman filter and smoothers for exponential family state space models.
URL <http://CRAN.R-project.org/package=KFAS>
- Hipel, K. W., Lennox, W. C., Unny, T. E., McLeod, A. I., 1975. Intervention analysis in water resources. *Water Resources Research* 11 (6), 855–861.
- Hipel, K. W., McLeod, A. I., 1994. *Time Series Modelling of Water Resources and Environmental Systems*. Elsevier, Amsterdam, electronic reprint available, <http://www.stats.uwo.ca/faculty/aim/1994Book/>.

- Hoffmann, T. J., 2011. Passing in command line arguments and parallel cluster/multicore batching in r with batch. *Journal of Statistical Software, Code Snippets* 39 (1), 1–11.
URL <http://www.jstatsoft.org/v39/c01>
- Hopwood, W. S., McKeown, J. C., Newbold, P., 1984. Time series forecasting models involving power transformations. *Journal of Forecasting* 3 (1), 57–61.
- Hornik, K., Leisch, F., 2001. Neural network models. In: Peña, D., Tiao, G. C., Tsay, R. S. (Eds.), *A Course in Time Series Analysis*. Vol. New York. Wiley, Ch. 13, pp. 348–364.
- Hothorn, T., Zeileis, A., Millo, G., Mitchell, D., 2010. lmtest: Testing Linear Regression Models. R package version 0.9-27.
URL <http://CRAN.R-project.org/package=lmtest>
- Hyndman, R. J., 2010. forecast: Forecasting functions for time series. R package version 2.17.
URL <http://CRAN.R-project.org/package=forecast>
- Hyndman, R. J., Khandakar, Y., 2008. Automatic time series forecasting: The forecast package for R. *Journal of Statistical Software* 27 (3), 1–22.
URL <http://www.jstatsoft.org/v27/i03>
- Hyndman, R. J., Koehler, A. B., Ord, J. K., Snyder, R. D., 2008. *Forecasting with Exponential Smoothing: The State Space Approach*. Springer-Verlag, New York.
- Iacus, S. M., 2008. *Simulation and Inference for Stochastic Differential Equations: With R Examples*. Springer Science+Business Media, LLC, New York.
- Iacus, S. M., 2009. sde: Simulation and Inference for Stochastic Differential Equations. R package version 2.0.10.
URL <http://CRAN.R-project.org/package=sde>
- Johansen, S., 1995. *Likelihood-Based Inference in Cointegrated Vector Autoregressive Models*. Oxford, Oxford.

- Kajitani, Y., Hipel, K. W., McLeod, A. I., 2005. Forecasting nonlinear time series with feed-forward neural networks: A case study of canadian lynx data. *Journal of Forecasting* 24, 105–117.
- Kedem, B., Fokianos, K., 2002. *Regression Models for Time Series Analysis*. Wiley, New York.
- Keenan, D. M., 1985. A Tukey nonadditivity-type test for time series nonlinearity. *Biometrika* 72, 39–44.
- Kleiber, C., Zeileis, A., 2008. *Applied Econometrics with R*. Springer, New York.
- Kwiatkowski, D., Phillips, P. C. B., Schmidt, P., Shin, Y., 1992. Testing the null hypothesis of stationarity against the alternative of a unit root: How sure are we that economic time series have a unit root? *Journal of Econometrics* 54, 159–178.
- Lee, T. H., White, H., Granger, C. W. J., 1993. Testing for neglected nonlinearity in time series models. *Journal of Econometrics* 56, 269–290.
- Leisch, F., 2002. Dynamic generation of statistical reports using literate data analysis. In: Härdle, W., Rönz, B. (Eds.), *COMPSTAT 2002 – Proceedings in Computational Statistics*. Physica-Verlag, Heidelberg, pp. 575–580.
- Leisch, F., 2003. Sweave and beyond: Computations on text documents. In: Hornik, K., Leisch, F., Zeileis, A. (Eds.), *Proceedings of the 3rd International Workshop on Distributed Statistical Computing*, Vienna, Austria. ISSN 1609-395X.
URL
<http://www.ci.tuwien.ac.at/Conferences/DSC-2003/Proceedings/>
- Lewis, P. A. W., Stevens, J. G., 1991. Nonlinear modeling of time series using multivariate adaptive regression splines (mars). *Journal of the American Statistical Association* 86 (416), 864–877.
- Li, W. K., 1994. Time series models based on generalized linear models: Some further results. *Biometrics* 50 (2), 506–511.

- Luethi, D., Erb, P., Oztiger, S., 2010. FKF: Fast Kalman Filter.
URL <http://CRAN.R-project.org/package=FKF>
- Lupi, C., 2011. CADFtest: Hansen's Covariate-Augmented Dickey-Fuller Test. R package version 0.3-1.
URL <http://CRAN.R-project.org/package=CADFtest>
- Lütkepohl, H., 2005. New Introduction to Multiple Time Series Analysis. Springer-Verlag, New York.
- Lütkepohl, H., Krätzig, M. (Eds.), 2004. Applied Time Series Econometrics. Cambridge University Press, Cambridge.
- MacKinnon, J. G., 1996. Numerical distribution functions for unit root and cointegration tests. *Journal of Applied Econometrics* 11 (601-618).
- McLeod, A., 1994. Diagnostic checking periodic autoregression models with application. *The Journal of Time Series Analysis* 15, 221–223, addendum, *Journal of Time Series Analysis* 16, p.647-648.
- McLeod, A., 1998. Hyperbolic decay time series. *The Journal of Time Series Analysis* 19, 473–484.
URL <http://www.jstatsoft.org/v23/i05>
- McLeod, A., Balcilar, M., 2011. pear: Package for Periodic Autoregression Analysis. R package version 1.2.
URL <http://CRAN.R-project.org/package=pear>
- McLeod, A., Yu, H., Krougly, Z., 2007. Algorithms for linear time series analysis: With R package. *Journal of Statistical Software* 23 (5), 1–26.
URL <http://www.jstatsoft.org/v23/i05>
- McLeod, A., Yu, H., Krougly, Z., 2011a. FGN: Fractional Gaussian Noise, estimation and simulation. R package version 1.4.
URL <http://CRAN.R-project.org/package=ltsa>
- McLeod, A., Zhang, Y., 2008a. Faster arma maximum likelihood estimation. *Computational Statistics and Data Analysis* 52 (4), 2166–2176.
- McLeod, A., Zhang, Y., 2008b. Improved subset autoregression: With R package. *Journal of Statistical Software* 28 (2), 1–28.
URL <http://www.jstatsoft.org/v28/i02>

- McLeod, A., Zhang, Y., Xu, C., 2011b. FitAR: Subset AR Model Fitting. R package version 1.92.
URL <http://CRAN.R-project.org/package=FitAR>
- McLeod, A. I., 2010. FitARMA: Fit ARMA or ARIMA Using Fast MLE Algorithm. R package version 1.4.
URL <http://CRAN.R-project.org/package=FitARMA>
- McLeod, A. I., Li, W. K., 1983. Diagnostic checking arma time series models using squared-residual autocorrelations. *Journal of Time Series Analysis* 4, 269–273.
- McLeod, A. I., Zhang, Y., 2006. Partial autocorrelation parameterization for subset autoregression. *Journal of Time Series Analysis* 27 (4), 599–612.
- Meyer, D., June 2002. Naive time series forecasting methods: The holt-winters method in package ts. *R News* 2 (2), 7–10.
- Milborrow, S., 2011. earth: Multivariate Adaptive Regression Spline Models. R package version 2.6-2.
URL <http://CRAN.R-project.org/package=earth>
- Moore, D. S., 2007. *The Basic Practice of Statistics*, 4th Edition. W. H. Freeman & Co., New York.
- Murrell, P., 2011. *R Graphics*, 2nd Edition. Chapman and Hall/CRC, Boca Raton.
- Nason, G., 2008. *Wavelet Methods in Statistics with R*. Springer-Verlag, New York.
- Nason, G., 2010. wavethresh: Wavelets statistics and transforms. R package version 4.5.
URL <http://CRAN.R-project.org/package=wavethresh>
- Peng, R., 2008. A method for visualizing multivariate time series data. *Journal of Statistical Software* 25 (Code Snippet 1), 1–17.
URL <http://www.jstatsoft.org/v25/c01>
- Percival, D. B., Walden, A. T., 1993. *Spectral Analysis For Physical Applications*. Cambridge University Press, Cambridge.

- Percival, D. B., Walden, A. T., 2000. Wavelet Methods for Time Series Analysis. Cambridge University Press, Cambridge.
- Petris, G., 2010. dlm: Bayesian and Likelihood Analysis of Dynamic Linear Models.
URL <http://CRAN.R-project.org/package=dlm>
- Petris, G., Petrone, S., Campagnoli, P., 2009. Dynamic Linear Models with R. Springer Science+Business Media, LLC, New York.
- Pfaff, B., 2006. Analysis of Integrated and Cointegrated Time Series with R. Springer, New York.
- Pfaff, B., 2008. Var, svar and svec models: Implementation within R package vars. Journal of Statistical Software 27 (4), 1–32.
URL <http://www.jstatsoft.org/v27/i04>
- Pfaff, B., 2010a. urca: Unit root and cointegration tests for time series data. R package version 1.2-5.
URL <http://CRAN.R-project.org/package=urca>
- Pfaff, B., 2010b. vars: VAR Modelling. R package version 1.4-8.
URL <http://CRAN.R-project.org/package=vars>
- Phillips, P. C. B., Ouliaris, S., 1990. Asymptotic properties of residual based tests for cointegration. Econometrica 58, 165–193.
- Phillips, P. C. B., Perron, P., 1988. Testing for a unit root in time series regression. Biometrika 75 (2), 335–346.
- R Development Core Team, 2011. Writing R Extensions. R Foundation for Statistical Computing, Vienna, Austria.
URL <http://www.R-project.org/>
- Revolution Computing, 2011. foreach: For each looping construct for R. R package version 1.3.2.
URL <http://CRAN.R-project.org/package=foreach>
- Ripley, B., 2011. nnet: Feed-forward Neural Networks and Multinomial Log-Linear Models. R package version 7.3-1.
URL <http://CRAN.R-project.org/package=nnet>

- Ripley, B. D., 1996. Pattern Recognition and Neural Networks. Cambridge University Press, New York.
- Ripley, B. D., June 2002. Time series in R 1.5.0. R News 2 (2), 2–7.
- Ritz, C., Streibig, J. C., 2008. Nonlinear Regression with R. Springer Science+Business Media, LLC, New York.
- Said, S. E., Dickey, D. A., 1984. Test for unit roots in autoregressive-moving average models of unknown order. Biometrika 71 (3), 599–607.
- Sarkar, D., 2008. Lattice: Multivariate Data Visualization with R. Springer, New York.
- Schmidberger, M., Morgan, M., Eddelbuettel, D., Yu, H., Tierney, L., Mansmann, U., 8 2009. State of the art in parallel computing with R. Journal of Statistical Software 31 (1), 1–27.
URL <http://www.jstatsoft.org/v31/i01>
- Shadish, W. R., Cook, T. D., Campbell, D. T., 2001. Experimental and Quasi-Experimental Designs for Generalized Causal Inference, 2nd Edition. Houghton Mifflin.
- Shumway, R. H., Stoffer, D. S., 2011. Time Series Analysis and Its Applications With R Examples, 3rd Edition. Springer.
- Smith, B., Field, C., 2001. Symbolic cumulant calculations for frequency domain time series. Statistics and Computing 11, 75–82.
- Spector, P., 2008. Data Manipulation with R. Springer-Verlag, Berlin.
- Teraesvirta, T., Lin, C. F., Granger, C. W. J., 1993. Power of the neural network linearity test. Journal of Time Series Analysis 14, 209–220.
- Tesfaye, Y. G., Anderson, P. L., Meerschaert, M. M., 2011. Asymptotic results for fourier-parma time series. Journal of Time Series Analysis 32 (2), 157–174.
- Thompson, M. E., McLeod, A. I., June 1976. The effects of economic variables upon the demand for cigarettes in Canada. The Mathematical Scientist 1, 121–132.

- Trapletti, A., 2011. tseries: Time series analysis and computational finance. R package version 0.10-25.
URL <http://CRAN.R-project.org/package=tseries>
- Tsay, R. S., 2010. Analysis of Financial Time Series, 3rd Edition. Wiley, New York.
- Tusell, F., 2011. Kalman filtering in R. Journal of Statistical Software 39 (2).
URL <http://www.jstatsoft.org/v39/i02>
- Ursu, E., Duchesne, P., 2009. On modelling and diagnostic checking of vector periodic autoregressive time series models. Journal of Time Series Analysis 30 (1), 70–96.
- Venables, W. N., Ripley, B. D., 2000. S Programming. Springer.
- Venables, W. N., Ripley, B. D., 2002. Modern Applied Statistics with S, 4th Edition. Springer.
- Vingilis, E., McLeod, A. I., Seeley, J., Mann, R. E., Stoduto, G., Compton, C., Beirness, D., 2005. Road safety impact of extended drinking hours in ontario. Accident Analysis and Prevention 37, 547–556.
- Whitcher, B., 2010. waveslim: Basic wavelet routines for one-, two- and three-dimensional signal processing. R package version 1.6.4.
URL <http://CRAN.R-project.org/package=waveslim>
- Wickham, H., 2009. ggplot2: Elegant graphics for data analysis. Springer New York.
URL <http://had.co.nz/ggplot2/book>
- Wilkinson, L., 1999. The Grammar of Graphics. Springer, New York.
- Wolfram Research, I., 2011. Mathematica Edition: Version 8.0. Champaign, Illinois.
- Wood, S., 2006. Generalized Additive Models: An Introduction with R. Chapman and Hall/CRC.

- Wuertz, D., 2010. fBasics: Rmetrics - Markets and Basic Statistics. R package version 2110.79.
URL <http://CRAN.R-project.org/package=fBasics>
- Wuertz, D., Chalabi, Y., 2011. timeSeries: Rmetrics - Financial Time Series Objects. R package version 2130.92.
URL <http://CRAN.R-project.org/package=timeSeries>
- Wuertz, D., et al., 2009a. fGarch: Rmetrics - Autoregressive Conditional Heteroskedastic Modelling. R package version 2110.80.
URL <http://CRAN.R-project.org/package=fGarch>
- Wuertz, D., et al., 2009b. fUnitRoots: Trends and Unit Roots. R package version 2100.76.
URL <http://CRAN.R-project.org/package=fUnitRoots>
- Würtz, D., 2004. Rmetrics: An Environment for Teaching Financial Engineering and Computational Finance with R. Rmetrics, ITP, ETH Zürich, Swiss Federal Institute of Technology, ETH Zürich, Switzerland.
URL <http://www.rmetrics.org>
- Zeileis, A., 2010. dynlm: Dynamic Linear Regression. R package version 0.3-0.
URL <http://CRAN.R-project.org/package=dynlm>
- Zeileis, A., Leisch, F., Hansen, B., Hornik, K., Kleiber, C., 2010. strucchange: Testing, Monitoring and Dating Structural Changes. R package version 1.4-4.
URL <http://CRAN.R-project.org/package=strucchange>
- Zeileis, A., Leisch, F., Hornik, K., Kleiber, C., 2002. Strucchange: An R package for testing for structural change in linear regression models. Journal of Statistical Software 7 (2), 1–38.
URL <http://www.jstatsoft.org/v07/i02>
- Zhang, Y., McLeod, A. I., 2006. Computer algebra derivation of the bias of burg estimators. Journal of Time Series Analysis 27, 157–165.
- Zhou, L., Braun, W. J., 2010. Fun with the r grid package. Journal of Statistics Education 18.
URL <http://www.amstat.org/publications/jse/v18n3/zhou.pdf>

- Zivot, E., Wang, J., 2006. Modeling Financial Time Series with S-PLUS, 2nd Edition. Springer Science+Business Media, Inc, New York.
- Zucchini, W., MacDonald, I. L., 2009. Hidden Markov Models for Time Series: A Practical Introduction using R, 2nd Edition. Chapman & Hall/CRC, Boca Raton.
- Zuur, A. F., Ieno, E. N., Meesters, E., 2009. A Beginner's Guide to R. Springer Science+Business Media, LLC, New York.