

# MS-E2148 Dynamic optimization

## Lecture 2

# Contents

- ▶ Transversality conditions in calculus of variations
- ▶ Solutions with corners
- ▶ Material Kirk 4

# Recap

- ▶ We derived Euler equation for the basic problem using the fundamental theorem of calculus of variations

# Calculus of variations

## Basic problem

- ▶ Let us find curve  $x(t)$  on a closed interval  $[t_0, t_f]$  that satisfies the boundary conditions  $x(t_0) = x_0$  and  $x(t_f) = x_f$  and is local extremum for the functional

$$J(x) = \int_{t_0}^{t_f} g(x, \dot{x}, t) dt \quad (1)$$

- ▶ This  $x = x^*$  is called the extremal
- ▶ The necessary condition for  $x^*$  is the Euler equation

$$g_x(x^*, \dot{x}^*, t) - \frac{d}{dt} g_{\dot{x}}(x^*, \dot{x}^*, t) = 0 \quad \forall t \in [t_0, t_f] \quad (2)$$

# Transversality conditions



## Different problems

- ▶ The basic problem is a two-point boundary value problem
- ▶ In practical problems the other end point can usually be free, when we get the following problems
  - ▶ Free final state
  - ▶ Free final time
  - ▶ Final state and time are independent
  - ▶ Final state and time are free but depend on each other
- ▶ Analysis is similar if we have free initial state/time
- ▶ The Euler equation is always one of the necessary conditions for the extremal independent of the boundary conditions!

# Transversality conditions

## Free final state

- ▶ Let us find the extremal for the functional (1) so that  $t_0$  and  $t_f$  are fixed and  $x(t_0) = x_0$ , but  $x(t_f)$  is free



# Transversality conditions

## Free final state

- ▶ Variation of  $J$  is

$$\begin{aligned}\delta J(x, \delta x) &= g_{\dot{x}}(x, \dot{x}, t)\delta x \Big|_{t_0}^{t_f} \\ &+ \int_{t_0}^{t_f} (g_x(x, \dot{x}, t) - \frac{d}{dt}g_{\dot{x}}(x, \dot{x}, t))\delta x dt\end{aligned}$$

- ▶ Now, the first term is not zero, since  $\delta x(t_f)$  is arbitrary

# Transversality conditions

Free final state

- ▶ Beside Euler, we have condition:

$$g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f) = 0 \quad (3)$$



# Transversality conditions

## Free final time

- ▶ Let us find extremal for the functional (1) so that  $t_0$  is fixed,  $x(t_0) = x_0$ , ja  $x(t_f) = x_f$ , but  $t_f$  is free
- ▶ Now beside the integration constants, we need to find what is the final time  $t_f$  on the extremal, since it is independent variable (not like  $x_f$  that depended on fixed  $t_f$ )

# Transversality conditions

## Free final time

- ▶ All extremal candidates end up on the horizontal line where  $x = x_f$ ; two curves in comparison end up on  $(x_f, t_f)$  and  $(x_f, t_f + \delta t_f)$
- ▶ The increment of the functional is

$$\begin{aligned}\Delta J &= \int_{t_0}^{t_f + \delta t_f} g(x, \dot{x}, t) dt - \int_{t_0}^{t_f} g(x^*, \dot{x}^*, t) dt \\ &= \int_{t_0}^{t_f} (g(x^* + \delta x, \dot{x}^* + \delta \dot{x}, t) - g(x^*, \dot{x}^*, t)) dt \\ &\quad + \int_{t_f}^{t_f + \delta t_f} g(x, \dot{x}, t) dt\end{aligned}$$

# Transversality conditions

## Free final time

- ▶ Let us expand the first term in the integral of  $\Delta J$  using Taylor series

$$\begin{aligned}\Delta J &= \int_{t_0}^{t_f} \left[ g_x(x^*, \dot{x}^*, t) \delta x + g_{\dot{x}}(x^*, \dot{x}^*, t) \delta \dot{x} \right] dt \\ &\quad + O(\delta x, \delta \dot{x}) + \int_{t_f}^{t_f + \delta t_f} g(x, \dot{x}, t) dt\end{aligned}$$

- ▶ The second term in the integral can be written as

$$\int_{t_f}^{t_f + \delta t_f} g(x, \dot{x}, t) dt = g(x(t_f), \dot{x}(t_f), t_f) \delta t_f + O(\delta t_f) \quad (4)$$

# Transversality conditions

## Free final time

- ▶ By partial integration and combining the term (4) to the increment:

$$\begin{aligned}\Delta J &= g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f)\delta x(t_f) + g(x(t_f), \dot{x}(t_f), t_f)\delta t_f \\ &\quad + \int_{t_0}^{t_f} \left[ g_x(x^*, \dot{x}^*, t) - \frac{d}{dt}g_{\dot{x}}(x^*, \dot{x}^*, t) \right] \delta x dt + O(\cdot)\end{aligned}$$

where we used that  $\delta x(t_0) = 0$ . Let us use the Taylor expansion on term  $g(x(t_f), \dot{x}(t_f), t_f)$ :

$$\begin{aligned}\Delta J &= g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f)\delta x(t_f) + g(x^*(t_f), \dot{x}^*(t_f), t_f)\delta t_f \\ &\quad + \int_{t_0}^{t_f} \left[ g_x(x^*, \dot{x}^*, t) - \frac{d}{dt}g_{\dot{x}}(x^*, \dot{x}^*, t) \right] \delta x dt + O(\cdot)\end{aligned}\tag{5}$$

# Transversality conditions

## Free final time

- ▶  $\delta x(t_f)$  depends on  $\delta t_f$ , so we can approximate (see the figure in two slides with  $\delta x_f = 0$ ):

$$\delta x(t_f) \approx -\dot{x}^*(t_f)\delta t_f \quad (6)$$

- ▶ The variation is solved from the increment (5) :

$$\begin{aligned} \delta J(x^*, \delta x) &= \left( (-g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f))\dot{x}^*(t_f) \right. \\ &\quad \left. + g(x^*(t_f), \dot{x}^*(t_f), t_f) \right) \delta t_f \\ &\quad + \int_{t_0}^{t_f} \left[ g_x(x^*, \dot{x}^*, t) - \frac{d}{dt} g_{\dot{x}}(x^*, \dot{x}^*, t) \right] \delta x dt \end{aligned}$$

# Transversality conditions

## Free final time

- ▶ Beside Euler, we have the transversality condition:


$$g(x^*(t_f), \dot{x}^*(t_f), t_f) - g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f)\dot{x}^*(t_f) = 0 \quad (7)$$

- ▶ **E.g.:**  $J = \int_1^T (2x + \dot{x}^2/2)dt$ ,  $x(1) = 4$ ,  $x(T) = 4$ , and  $T > 1$
- ▶ Euler:  $\ddot{x}^* = 2 \Rightarrow x^* = t^2 + c_1t + c_2$
- ▶ Transversality condition:  $4x^*(T) - \dot{x}^{*2}(T) = 0$
- ▶ Boundary conditions:  $x^*(1) = 4 = 1 + c_1 + c_2$ ;  
 $x^*(T) = 4 = T^2 + c_1T + c_2$ ;  
 $4x^*(T) - \dot{x}^{*2}(T) = 0 = 4c_2 - c_1^2$
- ▶ Solving constants  $c_1, c_2, T$  (3 constants, 3 equations) we get  $x^* = t^2 - 6t + 9$  and  $T = 5$

# Transversality conditions

## Free final state and time

- ▶ Let us find the extremal for the functional (1) so that  $t_0$  is fixed,  $x(t_0) = x_0$ , but  $x(t_f)$  and  $t_f$  are free



We get:  $\delta x_f \approx \delta x(t_f) + \dot{x}^*(t_f)\delta t_f$

# Transversality conditions

## Free final state and time

- ▶ Let us use the increment in (5), and use the above approximation

$$\begin{aligned}\delta J(x^*, \delta x) &= g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f) \delta x_f \\ &+ \left[ g(x^*(t_f), \dot{x}^*(t_f), t_f) \right. \\ &\quad \left. - g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f) \cdot \dot{x}^*(t_f) \right] \delta t_f \\ &+ \int_{t_0}^{t_f} \left[ g_x(x^*, \dot{x}^*, t) - \frac{d}{dt} g_{\dot{x}}(x^*, \dot{x}^*, t) \right] \delta x dt\end{aligned}\tag{8}$$



# Transversality conditions

## Free final state and time

- ▶ Variation is zero if Euler holds and the multipliers of  $\delta x_f$  and  $\delta t_f$  are zero
- ▶ There are two cases:
  - 1)  $t_f$  and  $x(t_f)$  are independent, when we get the transversality conditions

$$\begin{aligned}g(x^*(t_f), \dot{x}^*(t_f), t_f) &= 0 \\g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f) &= 0\end{aligned}\tag{9}$$

- 2)  $t_f$  depends on  $x(t_f)$ : the final point is on some curve  $x(t_f) = \theta(t_f)$ , and  $\delta x_f \approx \dot{\theta}(t_f)\delta t_f$  with transversality condition:

$$g_{\dot{x}}(x^*(t_f), \dot{x}^*(t_f), t_f) \left[ \dot{\theta}(t_f) - \dot{x}^*(t_f) \right] + g(x^*(t_f), \dot{x}^*(t_f), t_f) = 0\tag{10}$$

# Transversality conditions

## Free final state and time


- ▶ **E.g.:** find the shortest route from the origin to the curve/line  $\theta(t) = -5t + 15$
- ▶ We minimize  $\int_0^{t_f} \sqrt{1 + \dot{x}^2} dt$  with  $x(0) = 0$  and  $x(t_f) = -5t_f + 15$
- ▶ Euler:  $\ddot{x} = 0 \Rightarrow x^* = c_1 t + c_2$
- ▶ Boundary condition  $x(0) = 0 \Rightarrow c_2 = 0$
- ▶ Transversality condition:

$$\begin{aligned} 0 &= \frac{\dot{x}^*(t_f)}{\sqrt{1 + \dot{x}^{*2}(t_f)}} (-5 - \dot{x}^*(t_f)) + \sqrt{1 + \dot{x}^*(t_f)} \\ &= -5\dot{x}^*(t_f) + 1 \end{aligned}$$

- ▶ We get from the transversality condition and the solution candidate  $c_1 = 1/5$
- ▶ Final condition gives  $t_f = 5 \cdot (-5t_f + 15) \Rightarrow t_f = 75/26$

# Calculus of variations: solutions with corners

- ▶ In the basic problem, the extremal candidates are continuous and continuously differentiable (*smooth*); this is a strong restriction
- ▶ Now, we allow extremals that have piecewise continuous first time derivatives, e.g., the  $\dot{x}$  is continuous, except for a finite number of points on the interval  $[t_0, t_f]$
- ▶ Where  $\dot{x}$  is discontinuous, we say that  $x$  has a *corner*



# Calculus of variations: solutions with corners

## Weierstrass-Erdmann corner point conditions

- ▶ Let us assume that  $g$  has continuous first and second-order partial differentials with respect to all of its arguments on functional (1) and that  $t_0, t_f, x(t_0), x(t_f)$  are fixed
- ▶ We assume that  $\dot{x}$  has a point of discontinuity (corner) in some point  $t_1 \in (t_0, t_f)$  which is not known in advance
- ▶ The functional can be represented as

$$\begin{aligned} J(x) &= \int_{t_0}^{t_1} g(x, \dot{x}, t) dt + \int_{t_1}^{t_f} g(x, \dot{x}, t) dt \\ &\equiv J_1(x) + J_2(x) \end{aligned} \quad (11)$$

- ▶ We know that if  $x^*$  is extremal for  $J$ , then  $x^*(t)|_{t \in [t_0, t_1]}$  is extremal for  $J_1$  and  $x^*(t)|_{t \in [t_1, t_f]}$  is extremal for  $J_2$

# Calculus of variations: solutions with corners

## Weierstrass-Erdmann conditions

- ▶ Let us denote  $t_1^-$  and  $t_1^+$  as the left and right-hand side of the point of discontinuity
- ▶ Since the corner point coordinates are free, the variation  $\delta J(x^*, \delta x) =$

$$\begin{aligned} & g_{\dot{x}}(x^*(t_1^-), \dot{x}^*(t_1^-), t_1^-) \delta x_1 \\ & + \left[ g(x^*(t_1^-), \dot{x}^*(t_1^-), t_1^-) - g_{\dot{x}}(x^*(t_1^-), \dot{x}^*(t_1^-), t_1^-) \dot{x}^*(t_1^-) \right] \delta t_1 \\ & + \int_{t_0}^{t_1} \left[ g_x(x^*, \dot{x}^*, t) - \frac{d}{dt} g_{\dot{x}}(x^*, \dot{x}^*, t) \right] \delta x dt \\ & - g_{\dot{x}}(x^*(t_1^+), \dot{x}^*(t_1^+), t_1^+) \delta x_1 \\ & - \left[ g(x^*(t_1^+), \dot{x}^*(t_1^+), t_1^+) - g_{\dot{x}}(x^*(t_1^+), \dot{x}^*(t_1^+), t_1^+) \dot{x}^*(t_1^+) \right] \delta t_1 \\ & + \int_{t_1}^{t_f} \left[ g_x(x^*, \dot{x}^*, t) - \frac{d}{dt} g_{\dot{x}}(x^*, \dot{x}^*, t) \right] \delta x dt \end{aligned} \tag{12}$$

# Calculus of variations: solutions with corners

## Weierstrass-Erdmann conditions

- ▶ Since  $\delta x_1$  and  $\delta t_1$  are arbitrary, the necessary conditions beside Euler for vanishing variation are

$$g_{\dot{x}}(x^*(t_1^-), \dot{x}^*(t_1^-), t_1^-) = g_{\dot{x}}(x^*(t_1^+), \dot{x}^*(t_1^+), t_1^+)$$

and

$$\begin{aligned} & g(x^*(t_1^-), \dot{x}^*(t_1^-), t_1^-) - \left[ g_{\dot{x}}(x^*(t_1^-), \dot{x}^*(t_1^-), t_1^-) \right] \dot{x}^*(t_1^-) \\ &= g(x^*(t_1^+), \dot{x}^*(t_1^+), t_1^+) - \left[ g_{\dot{x}}(x^*(t_1^+), \dot{x}^*(t_1^+), t_1^+) \right] \dot{x}^*(t_1^+) \end{aligned}$$

- ▶  $\Rightarrow$  functions  $g_{\dot{x}}$  and  $g - g_{\dot{x}}\dot{x}$  must be continuous over the corner

# Calculus of variations: solutions with corners

## Example 1

- ▶  $J(x) = \int_0^{\pi/2} [\dot{x}^2 - x^2] dt$ ,  $x(0) = 0$  and  $x(\pi/2) = 1$
- ▶ Euler:  $\ddot{x} + x = 0 \Rightarrow x^* = c_3 \cos t + c_4 \sin t$
- ▶ W-E corner conditions:

$$2\dot{x}^*(t_1^-) = 2\dot{x}^*(t_1^+)$$

and

$$\begin{aligned} & \dot{x}^{*2}(t_1^-) - x^{*2}(t_1^-) - 2\dot{x}^*(t_1^-)\dot{x}^*(t_1^-) \\ &= \dot{x}^{*2}(t_1^+) - x^{*2}(t_1^+) - 2\dot{x}^*(t_1^+)\dot{x}^*(t_1^+) \end{aligned}$$

which require that  $\dot{x}$  is continuous over  $t_1$ ; i.e.,  $x^*$  *cannot have a corner*

# Calculus of variations: solutions with corners

## Example 2

- ▶  $J(x) = \int_0^2 [\dot{x}^2 - 1]^2 dt$ ,  $x(0) = 0$ ,  $x(2) = 0$
- ▶ Euler:  $\dot{x}^3 - \dot{x} = c_1 \Rightarrow x^* = c_2 t + c_3$
- ▶ Boundary conditions:  $x^* = 0$ , where  $J(x^*) = 2$
- ▶ Corner conditions:

$$\begin{aligned} 4\dot{x}^*(t_1^-)[\dot{x}^{*2}(t_1^-) - 1] &= 4\dot{x}^*(t_1^+)[\dot{x}^{*2}(t_1^+) - 1] \\ -(3\dot{x}^{*2}(t_1^-) + 1)[\dot{x}^{*2}(t_1^-) - 1]^2 &= -(3\dot{x}^{*2}(t_1^+) + 1)[\dot{x}^{*2}(t_1^+) - 1]^2 \end{aligned}$$

- ▶ The first equation is satisfied when  $\dot{x}^*(t_1^-) = -1, 0, 1$  and  $\dot{x}^*(t_1^+) = -1, 0, 1$
- ▶ The second is satisfied when  $\dot{x}^*(t_1^-) = -1, 1$  and  $\dot{x}^*(t_1^+) = -1, 1$



# Calculus of variations: solutions with corners

## Example 2

- ▶ In the solution, we either have  $\dot{x}^*(t_1^-) = 1$  and  $\dot{x}^*(t_1^+) = -1$  or  $\dot{x}^*(t_1^-) = -1$  and  $\dot{x}^*(t_1^+) = 1$
- ▶ Putting them together

$$\begin{cases} x^* = t, & t \leq t_1 = 1 \\ x^* = -t + 2, & t \geq t_1 = 1 \end{cases}$$

- ▶ This gives solution with  $J(x^*) = 0$  which is surely the global minimum (check the integrand)

# Summary

- ▶ On the extremal, Euler equation must hold
- ▶ ... and it is complemented with transversality conditions and Weierstrass-Erdmann corner conditions
- ▶ Transversality and corner conditions can be generalized to vector-valued functions