

MS-C1620 Statistical inference

4 Inference for binary data

Joni Virta

Department of Mathematics and Systems Analysis
School of Science
Aalto University

Academic year 2018–2019
Period III - IV

Contents

- 1 Binary data
- 2 Single binary sample
- 3 Two binary samples
- 4 Lecture quiz

Binary observations

In many applications the observations are binary.

- Something is true/false.
- Something happened/did not happen.
- Someone belongs/does not belong to a group.

In such a case the observations are most conveniently coded as 0/1.

Recall that if we have a iid sample of binary observations, their distribution is necessarily the *Bernoulli distribution*.

Bernoulli distribution

The random variable x is said to obey the Bernoulli distribution with the probability of success p if,

$$\mathbb{P}(x = 1) = p \quad \text{and} \quad \mathbb{P}(x = 0) = 1 - p.$$

The expected value and variance of x are,

$$\begin{aligned}\mathbb{E}(x) &= p \\ \text{Var}(x) &= p(1 - p).\end{aligned}$$

That is, the Bernoulli distribution has only a single parameter to estimate.

The sum of n i.i.d. Bernoulli random variables with the success probability p has the binomial distribution with the parameters n and p .

Contents

- 1 Binary data
- 2 Single binary sample**
- 3 Two binary samples
- 4 Lecture quiz

Approximate confidence interval

Central limit theorem can be used to obtain a confidence interval for the success probability p of a Bernoulli distribution.

Let x_1, x_2, \dots, x_n be an i.i.d. sample from the Bernoulli distribution with the success probability/expected value p .

For large n , a level $100(1 - \alpha)\%$ confidence interval for the success probability p is obtained as

$$\left(\hat{p} - z_{\alpha/2} \frac{\sqrt{\hat{p}(1 - \hat{p})}}{\sqrt{n}}, \hat{p} + z_{\alpha/2} \frac{\sqrt{\hat{p}(1 - \hat{p})}}{\sqrt{n}} \right),$$

where \hat{p} is the observed proportion of successes and $z_{\alpha/2}$ is the $(1 - \alpha/2)$ -quantile of the standard normal distribution.

One-sample proportion test

To test whether the success probability of a Bernoulli distribution equals some pre-specified value, we employ **one-sample proportion test**.

One-sample proportion test, assumptions

Let x_1, x_2, \dots, x_n be an i.i.d. sample from a Bernoulli distribution with the success probability p .

One-sample proportion test, hypotheses

$$H_0 : p = p_0 \quad H_1 : p \neq p_0.$$

One-sample proportion test

One-sample proportion test, test statistic

- The test statistic,

$$C = \sum_{i=1}^n x_i,$$

follows the binomial distribution with parameters n and p_0 under H_0 .

- Under H_0 , the test statistic has $E[C] = np_0$ and $\text{Var}(C) = np_0(1 - p_0)$ and both large and both **large** and **small** values of the test statistic suggest that the null hypothesis H_0 is false.

The distribution of the test statistic C is tabulated and statistical software calculate exact p -values of the test.

Asymptotic one-sample proportion test

If the sample size is large, then under the null hypothesis H_0 the standardized test statistic,

$$Z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}},$$

where \hat{p} is the unbiased estimator $\hat{p} = \frac{1}{n} \sum_{i=1}^n x_i$ of the parameter p , follows approximately the standard normal distribution.

The approximation is usually accurate enough if $n\hat{p} > 10$ and $n(1 - \hat{p}) > 10$. For smaller sample sizes one should rely on the exact distribution of the test statistic C .

Contents

- 1 Binary data
- 2 Single binary sample
- 3 Two binary samples**
- 4 Lecture quiz

Two-sample proportion test

The one-sample proportion test can be seen as the equivalent of t -test when the normal distribution is replaced by the Bernoulli distribution.

As with t -test, a two-sample version readily follows and in **two-sample proportion test** parameters of two independent Bernoulli-distributed samples are compared.

Two-sample proportion test, assumptions

Let x_1, x_2, \dots, x_n be an i.i.d. sample from a Bernoulli distribution with the success probability p_x and let y_1, y_2, \dots, y_m be an i.i.d. sample from a Bernoulli distribution with the success probability p_y . Furthermore, let the two samples be independent.

Two-sample proportion test, hypotheses

$$H_0 : p_x = p_y \quad H_1 : p_x \neq p_y.$$

Two-sample proportion test

One-sample proportion test, test statistic

- Calculate the sample proportions

$$\hat{p}_x = \frac{1}{n} \sum_{i=1}^n x_i, \quad \hat{p}_y = \frac{1}{m} \sum_{i=1}^m y_i, \quad \hat{p} = \frac{n\hat{p}_x + m\hat{p}_y}{n + m}.$$

- The test statistic,

$$Z = \frac{\hat{p}_x - \hat{p}_y}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n} + \frac{1}{m}\right)}},$$

follows *for large n* under H_0 the standard normal distribution.

- Both **large** and **small** values of the test statistic suggest that the null hypothesis H_0 is false.

The normal approximation is usually good enough if $n\hat{p}_x > 5$, $n(1 - \hat{p}_x) > 5$, $m\hat{p}_y > 5$ and $m(1 - \hat{p}_y) > 5$.

Frequency tables

Assuming a “paired binary sample”, the previous test is no longer valid.

id	X	Y
1	0	1
2	0	0
3	0	1
4	1	1
\vdots	\vdots	\vdots

This kind of data is most conveniently represented in a **frequency table**.

	Y = 0	Y = 1
X = 0	173	40
X = 1	65	53

Inference for frequency tables is discussed next time.

Contents

- 1 Binary data
- 2 Single binary sample
- 3 Two binary samples
- 4 Lecture quiz**

Lecture quiz

A lecture quiz to determine what you have learned thus far!

Answer the following questions on your own or in small groups.

Lecture quiz

Question 1

Consider the following random sample: 5, -4, -2, 2. Calculate the following sample quantities:

- 1 Sample mean
- 2 Sample standard deviation
- 3 Sample median
- 4 Sample median absolute deviation
- 5 Sample range
- 6 Signs of the sample points
- 7 Ranks of the sample points
- 8 Signed ranks of the sample points with respect to distance to 0.

Lecture quiz

Question 2

Give concrete examples when you would/would not use the following measures of location:

- 1 Sample mean
- 2 Sample median
- 3 Mode

Question 3

Give concrete examples when you would/would not use the following measures of scatter:

- 1 Standard deviation
- 2 Median absolute deviation
- 3 Sample range

Lecture quiz

Question 4

What does it mean in practice if:

- The confidence interval of a parameter is narrow
- The significance level of a test is set low
- The p -value of a test is high
- Type I error occurs in a statistical test
- Type II error occurs in a statistical test

Lecture quiz

Question 5

How would you visualize the following samples:

- The heights of the male and female students attending a course.
- The exam points (0-24) on a large course.
- The proportions of faulty products produced by 5 different production lines.
- Stock prices of 3 companies.
- The monthly salaries and postal codes of adults living in Helsinki area.

Lecture quiz

Question 6

The following plots show the distributions of the test statistics of t -test, sign test and signed rank test for the null hypothesis of zero location when the data is a sample of size $n = 10$ from the standard normal distribution. Which plot corresponds to which test?

