


Solid-solid and solid-liquid interphases


The modelling of solid interphases is rather difficult since the interphases atomic structure is mostly unknown and also the reorganization barriers in solid systems are very large. This means that the interphase is at best on a good local minima. On the other hand the grain boundaries are a very important for materials macroscopic properties.


Example: Si|SiO₂ interphase.



Grain boundaries


Every real material consists of grains and real materials are full of grain boundaries. We can assume that each grain is a single crystal in which the atoms are perfectly oriented. The grain size varies a lot but is usually on the micrometer scale. Typically the grains have irregular shapes. Now the grain boundaries can be classified with the tilt and twist angles





Schematic representations of a tilt boundary (top) and a twist boundary between two idealised grains.

Atoms in these ideal boundaries will relax to minimize the interaction energy. The simplest grain boundary is the one with only tilt. If the angle is small boundary can be described with a dislocation. There will an extra atomic layer in the systems and after some layer again a new layer. As one can imagine the interphase structure can be complex and on atomic level they have not been modelled much. The grain boundaries will also move (slowly) since the atoms will diffuse. Often this movement is not wanted. One can add some impurities (atoms) to prevent the boundaries to move. (Zener pinning)



There has been very little ab initio modelling of grain boundaries. I found one interesting article A.L-S.Chua et al. Nature Mat. 9 (2010), 418.

Abstract

Recent years have seen great advances in our ability to predict crystal structures from first principles. However, previous algorithms have focused on the prediction of bulk crystal structures, where the global minimum is the target. Here, we present a general atomistic approach to simulate in multicomponent systems the structures and free energies of grain boundaries and heterophase interfaces with fixed stoichiometric and non-stoichiometric compositions. The approach combines a new genetic algorithm using empirical interatomic potentials to explore the configurational phase space of boundaries, and thereafter refining structures and free energies with first-principles electronic structure methods. We introduce a structural order parameter to bias the genetic algorithm search away from the global minimum (which would be bulk crystal), while not favouring any particular structure types, unless they lower the energy. We demonstrate the power and efficiency of the algorithm by considering non-stoichiometric grain boundaries in a ternary oxide, SrTiO₃.

One the other hand there has been some publications using empirical models. A good example is: D.E. Spearot and D.L. McDowell, J. Eng. Mater. Technol. **131**, 041204 (2009).

Solid liquid interphase – water molecules on a surface

The real solid water interphase has not been studied much with ab initio methods. The reason is simple – the calculations are time consuming. On the other hand single and few water molecules has been studied quite a lot. The water molecule will bound to almost any surface. It mostly stable but on for example transition metal oxide surfaces (like TiO₂) the water molecules can dissociate. We look first the simple metal surfaces. Compared to many other molecules the water is more problematic since it form easily hydrogen bonds. Then the cluster structures depend both one the interaction with the surface and the hydrogen bonds.

A good article is: S.Meng, E.G.Wang and S.Gao, Phys. Rev. B 69, 195404 (2004)

With small amount of water the situation is simple. The water molecules are usually oxygen toward the metal. If there are more molecules they form hydrogen bonds. The O-O distance is seldom close to the M-M distance so the molecules in clusters are usually not on certain specific positions (like top, bridge or hollow).





Figure: water clusters on Pt(111) surface


But when there are more waters the over layer structure can be complex.



The surface cell is $\sqrt{39} \times \sqrt{39} R 16.1^\circ$. Note that the waters are not exactly on the Pt atoms and this is the main reason to the large unit cell. Also the water layer is smooth. There is quite a bit of variation in the height.

Liquid water solid interphase


As said earlier there are very few ab initio studies of liquid solid interphase. One of the studied systems is water-NaCl. The system is relatively simple. There is a slab of NaCl and on top of that some layers of water. The NaCl is naturally ionic so the water molecules will orient strongly near to the surface.



Lähde: L.-M. Liu M. Krack and A. Michaelides, *J. Chem. Phys.* 130, 234702, (2009) and J-C Chen, P. Spijker, B. Reischl, N. Holmberg, K. Laasonen and A. S. Foster, 'ab initio Kinetic Monte Carlo simulations of dissolution at the NaCl-water interface', *PCCP*, 16, 22545-22554, (2014).

Next there are some plot of vertical density profile with both empirical and ab initio simulations. The first layer is clear but it varies a lot from model to model. Note that in the first picture the DFT results are very noisy. In the second picture the DFT simulations are much longer and the systems is also larger. The second DFT simulations shows that the maximum density is rather low (ca. 1.6 g/cm³) compared to many of the empirical models (2.0-3.0 g/cm³)


Figure 1: Density profiles with several ion models near the NaCl surface. Figure (a) is with TIP3P water model and (b) with SPC/E model. From: L.-M. Liu M. Krack and A. Michaelides, *J. Chem. Phys.* 130, 234702, (2009)).





Liquid water on NaCl(100) surface

- PBE (4x4) cell
- PBE (3x3) cell
- BLYP (4x4) cell


Chen et al. PCCP (2014)



The structure of water is interesting but also the Na and Cl dissociation can be studied (Chen et al. PCCP (2014)). Computationally the atoms can be pulled from the surface and the force acting to the atom can be computed. This force is rather noisy so relatively long simulations are needed.




After the simulation one can compute the reaction barriers. Below the barrier have been tested with different equivalent sites.




Also different surface sites can be studied. We have studied flat surface, sites near vacancy, step sites and corner sites.

NaCl (100) surface models



The important observation is that the barriers are very different




It is rather obvious that the corner atoms dissociates easily and also the steps atoms dissolves much easier than surface atoms. We have also made a

Kinetic Monte Carlo model for this system. With the KCM model systems of thousands of atoms can be studied.


system	Barrier (eV)	System	Barrier (eV)
Flat Cl-	0.75	Corner Cl-	0.17
Flat Na+	0.78	Corner Na+	0.05
Step Cl-	0.23	Vacancy Cl-	0.27
Step Na+	0.26	Vacancy Na+	0.10

The KCM predict that cube of NaCl will become round. On the left there is the simulation result and on the right an experimental picture of the NaCl particle in humid atmosphere. The structure is same but the size is different (simulation 14 nm and exp 160 nm)

(Exp: A. Bruzewicz, A. Checco, B. M. Ocko, E. R. Lewis, R. L. McGraw and S. E. Schwartz, JCP 134 (2011) 2011)




D=14 nm



Also TiO₂ - water interphase has been modelled


We study a collection of water molecules interacting with a perfect TiO₂(110) surface using *ab initio* molecular dynamics simulations. No water dissociation is observed. When we zoom in on the interface zone we see that water molecules can jump away from the first layer and go into the second layer. More details can be found in [Liu et al, Phys. Rev. B. 82, 161415\(R\) \(2010\)](#)





Electrochemical modelling

Another very interesting field of water solid interphase modelling is the modelling of electrochemical reactions. The reaction are relatively "easy", like hydrogen evolution


Hydrogen evolution reaction (HER)





Detailed reaction




(H^* = H bonded to catalyst)



We have done some modelling with Carbon Nanotubes (CNT) as catalyst. Naturally the CNT has to be in water. Then the hydrogen reaction need to be modelled. An extra problem but the key in electrochemistry is the voltage in the system. This makes the catalyst to be charged and this has a big effect to the reactivity. The charge is easy to include to the computations (but it is not easy to find the relation between the voltage and charge). The next step is to compute the barriers with some charged systems.



The Volmer reaction ($\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{H-CNT}$) and its activation energy. Note that the activation energy is a rather smooth but sensitive function of the potential. There are two systems: pure CNT and N-doped CNT (NCNT).



The Heyrovsky reaction ($\text{H}^+(\text{aq}) + \text{H-CNT} + \text{e}^- \rightarrow \text{H}_2 + \text{CNT}$) and its activation energy. Note that the Heyrovsky barrier is much higher, so the Heyrovsky reaction is the rate-limiting step. The potential range is realistic since around -1.7 V water starts breaking. The Heyrovsky reaction barrier is rather high, so CNT or NCNT are not good catalysts for HER. The best catalyst for HER is Pt and also that has been studied computationally.

In general

